ATTORNEY'S DOCKET NUMBER FORM PTO-1390 (Modified) REV 11-2000) U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE 29729/37581 TRANSMITTAL LETTER TO THE UNITED STATES U.S APPLICATION NO. (IF KNOWN, SEE 37 CFR DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371 INTERNATIONAL APPLICATION NO. INTERNATIONAL FILING DATE PRIORITY DATE CLAIMED PCT/AU00/00011 11 January 2000 11 January 1999 TITLE OF INVENTION CATALYTIC MOLECULES APPLICANT(S) FOR DO/EO/US DAVID G. ATKINS, ANDREW BAKER, LEVON M. KHACHIGIAN Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information: This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. 2. This is an express request to begin national examination procedures (35 U.S.C 371(f)). The submission must include itens (5), (6), (9) and (24) indicated below. The US has been elected by the expiration of 19 months from the priority date (Article 31). 4. A copy of the International Application as filed (35 U.S.C. 371 (c) (2)) is attached hereto (required only if not communicated by the International Bureau). b. □ has been communicated by the International Bureau. is not required, as the application was filed in the United States Receiving Office (RO/US). An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)). a. 🔲 is attached hereto. b. 🗆 has been previously submitted under 35 U.S.C. 154(d)(4). Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3)) are attached hereto (required only if not communicated by the International Bureau). have been communicated by the International Bureau. c. 🗆 have not been made; however, the time limit for making such amendments has NOT expired. have not been made and will not be made. 8. An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). 9 An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)). An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)). 10. Ø 11. A copy of the International Preliminary Examination Report (PCT/IPEA/409). 12. A copy of the International Search Report (PCT/ISA/210). Items 13 to 20 below concern document(s) or information included: An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 14. An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. \boxtimes 15. A FIRST preliminary amendment. 16. A SECOND or SUBSEQUENT preliminary amendment. 17. A substitute specification. 18. A change of power of attorney and/or address letter. 19. \boxtimes A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825. 20. \Box A second copy of the published international application under 35 U.S.C. 154(d)(4). 21 A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4). 22. \boxtimes Certificate of Mailing by Express Mail 23. Other items or information:

JC18 Rec'd PCT/PTO 1 1 JUL 2001

U.S. APPLICATION	9/889075	1	DOCKET NUMBER 19/37581				
24. The fol	lowing fees are submitted:.	CALCULATION	S PTO USE ONLY				
	L FEE (37 CFR 1.492 (a) (1) -	(5)):				CABCOCATION	S THOUSE CHE!
internationa	rnational preliminary examination I search fee (37 CFR 1.445(a)(2)) ional Search Report not prepared	paid to USPTO		\$100	0.00		
☐ Internationa USPTO but	l preliminary examination fee (37 International Search Report prepa	CFR 1.482) not paid to ared by the EPO or JPO		\$86	0.00		
International but international	l preliminary examination fee (37 onal search fee (37 CFR 1.445(a)	CFR 1.482) not paid to (2)) paid to USPTO	USPTC) . \$71	0.00	i :	
Internationa but all claim	I preliminary examination fee (37 is did not satisfy provisions of PC	CFR 1.482) paid to USI T Article 33(1)-(4)	PTO 	\$69	0.00		
	l preliminary examination fee (37 as satisfied provisions of PCT Art	icle 33(1)-(4)			0.00		
	ENTER APPROPRIA	ATE BASIC FEE	AMO	OUNT =		\$1,000.00	
	00 for furnishing the oath or decla liest claimed priority date (37 CI		□ 20) 🗆 30)	\$0.00	
CLAIMS	NUMBER FILED	NUMBER EXTR	A	RATE			
Total claims	43 - 20 =	23		x \$18.0	0	\$414.00	
Independent claims		0		x \$80.0	0	\$0.00	
Multiple Dependent	Claims (check if applicable).	ADOVE CAT C	TT 4.00	IONG		\$270.00	
- Annliant al-		ABOVE CALCU			=	\$1,684.00	
reduced by 1/2	ms small entity status. (See 37 CF	K 1.27). The fees indica	ted abov	ve are		\$0.00	·
			SUB	FOTAL	=	\$1,684.00	
Processing fee of \$1 nonths from the ear	30.00 for furnishing the English eliest claimed priority date (37 CF	translation later than FR 1.492 (f)).	□ 20) 🗆 30) +	\$0.00	
		TOTAL NATIO	ONAI	FEE	=	\$1,684.00	
Fee for recording th accompanied by an	e enclosed assignment (37 CFR 1 appropriate cover sheet (37 CFR)	.21(h)). The assignment 3.28, 3.31) (check if ap	t must b	e e).		\$0.00	
		TOTAL FEES E	CNCL	OSED	=	\$1,684.00	
•					i	Amount to be: refunded	\$
						charged	\$
a. 🗵 A ch	eck in the amount of\$1,684	to cover the ab	ove fees	is enclosed			
	se charge my Deposit Account No plicate copy of this sheet is enclo		the amo	ount of		to cover t	he above fees.
	Commissioner is hereby authorize eposit Account No. 13-2855					uired, or credit any o	overpayment
d. Fees	are to be charged to a credit card	. WARNING: Informati			•	•	
NOTE: Where an	appropriate time limit under 37	CFR 1.494 or 1.495 ha	s not b	een met, a p			
1.137(a) or (b)) mu	st be filed and granted to restor	e the application to pen	dingst	itus.	i	1 1, 11	A.
SEND ALL CORRE	ESPONDENCE TO-			\ac-	1. [do IMILana	16.
11/1/ T 1 1 1 1 0 1		Ì		SIGNATI	IRE/	1 V was	~
WILLIAMS, Jose MARSHALL, O'T		(' .)	./		\circ	
MURRAY & B	ORUN	1	'	\ 	. Wil	liams, Jr.	
6300 Sears Tower				NAME			
233 South Wacker Chicago, Illinois 6		İ		38,659			
(312) 474-6300				REGISTR	ATIO	N NUMBER	
		į		11 July 2	001		
				DATE			
		į.					



Patrio Co mod 1 Up - 200

PATENT 30397/37581

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Atkins et al. Serial No.: 09/889,075 U.S. National Phase of PCT/AU00/00011 Filed: January 11, 2000 For: Catalytic Molecules	 I hereby certify that this paper is being deposited with the United States Postal Service with sufficient postage as first class mail in an envelope addressed to: Commissioner for Patents, Washington, D.C. 20231 on this date:
Group Art Unit: Not Yet Assigned) September 4, 2002
Examiner: Not Yet Assigned	David A. Gass Registration No. 38,153 Attorney for Applicants

PRELIMINARY AMENDMENT

Commissioner for Patents Washington, DC 20231

Sir:

Please amend the above-identified application as follows, prior to calculation of the filing fee (with extra claims) and prior to examination of the merits:

IN THE CLAIMS:

AMEND Claims 48, 49 and 59as follows:

- 48. (AMENDED) A pharmaceutical composition comprising a DNAzyme according to claim 20 and a pharmaceutically acceptable carrier.
- 49. (AMENDED) A method of inhibiting EGR-1 activity in cells which comprises exposing the cell to a DNAzyme according to claim 20.

09/889,075

59. (AMENDED) An angioplastic stent for inhibition of the onset of restenosis, which comprises an angioplastic stent operably coated with a prophylactially effective dose of DNAzyme according to claim 20.

REMARKS

The applicant respectfully submits that all claims are now of proper form and scope for allowance. Early and favorable consideration is respectfully requested.

The foregoing amendments are intended solely to reduce the PTO fees for multiple dependent claims and claims in excess of 20. The applicants reserve the right to pursue subject matter of any cancelled claim in this application or any related (e.g., continuing) application.

Attached hereto is a marked-up version of the changes made to the claims by the foregoing amendments. The attached page is captioned "Version with markings to show changes made."

Respectfully submitted,

MARSHALL, GERSTEIN & BORUN

Bv:

David A. Gass Reg. No. 38,153 6300 Sears Tower 233 South Wacker Drive

Chicago, Illinois 60606-6402

(312) 474-6300

September 3, 2002

09/889,075

30397/37581

VERSION WITH MARKINGS TO SHOW CHANGES MADE

IN THE CLAIMS:

AMEND Claims 48, 49 and 59 as follows:

- 48. (AMENDED) A pharmaceutical composition comprising a DNAzyme according to [any one of claims 20 to 47] <u>claim 20</u> and a pharmaceutically acceptable carrier.
- 49. (AMENDED) A method of inhibiting EGR-1 activity in cells which comprises exposing the cell to a DNAzyme according to [any one of claims 20 to 47] <u>claim 20</u>.
- 59. (AMENDED) An angioplastic stent for inhibition of the onset of restenosis, which comprises an angioplastic stent operably coated with a prophylactially effective dose of DNAzyme according to [any one of claims 20 to 47] claim 20.

09/889075 JC18 Rec'd PCT/PTO 1 1 JUL 2001

PATENT 29729/37581

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Application of: David G. Atkins et al)	"EXPRESS MAIL" mailing label No.
)	EL564462123US
U.S. National Phase of)	
International Application No:)	I hereby certify that this paper is being
PCT/AU/00/00011)	deposited with the United States Postal
filed January 11, 2000)	Service as first class mail, postage
)	prepaid, in an envelope addressed to:
Serial No: Not yet assigned)	Box PCT, Commissioner for Patents,
•)	Washington, D.C. 20231, on this date
Filed: herewith)	
)	July 11, 2001
Title: CATALYTIC MOLECULES)	111111111111111111111111111111111111
)	
Group Art Unit: Not yet assigned)	Kichaul tr
)	Richard Zimmermann
Examiner: Not yet assigned)	

PRELIMINARY AMENDMENT

Box PCT Commissioner for Patents Washington, D.C. 20231

Dear Sir:

This amendment is being filed with the U.S. National filing of the above identified PCT application. The Applicant requests entry of this amendments prior to calculation of the filing fee and prior to examination on the merits.

AMENDMENTS

In the Abstract:

Please add to the application the abstract attached hereto.

In the claims:

Please cancel claims 1-19 and insert the following claims 20-62 as shown below:

- 20. A DNAzyme which specifically cleaves EGR-1 mRNA, the DNAzyme comprising (i) a catalytic domain which cleaves mRNA at a purine:pyrimidine cleavage site; (ii) a first binding domain continuous with the 5' end of the catalytic domain; and (iii) a second binding domain continuous with the 3' end of the catalytic domain, wherein the binding domains are sufficiently complementary to the two regions immediately flanking a purine:pyrimidine cleavage site within the region of EGR-1 mRNA corresponding to nucleotides 168-332 as shown in SEQ ID No: 1, such that the DNAzyme cleaves the EGR-1 mRNA.
- 21. A DNAzyme as claimed in claim 20 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
- 22. A DNAzyme as claimed in claim 20 in which the cleavage site is selected from the group consisting of
 - (i) the GU site corresponding to nucleotides 198-199;
 - (ii) the GU site corresponding to nucleotides 200-201;
 - (iii) the GU site corresponding to nucleotides 264-265;
 - (iv) the AU site corresponding to nucleotides 271-272;
 - (v) the AU site corresponding to nucleotides 301-302:
 - (vi) the GU site corresponding to nucleotides 303-304; and
 - (vii) the AU site corresponding to nucleotides 316-317.
- 23. A DNAzyme as claimed in claim 22 in which the cleavage site is the AU site corresponding to nucleotides 271-272.
- 24. A DNAzyme as claimed in claim 22 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
- 25. A DNAzyme as claimed in claim 23 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
- 26. A DNAzyme as claimed in claim 20 in which the catalytic domain has the nucleotide sequence GGCTAGCTACAACGA [SEQ. ID. NO: 2].
- 27. A DNAzyme as claimed in claim 26 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.

- 28. A DNAzyme as claimed in claim 26 in which the cleavage site is selected from the group consisting of
 - (i) the GU site corresponding to nucleotides 198-199;
 - (ii) the GU site corresponding to nucleotides 200-201;
 - (iii) the GU site corresponding to nucleotides 264-265;
 - (iv) the AU site corresponding to nucleotides 271-272;
 - (v) the AU site corresponding to nucleotides 301-302;
 - (vi) the GU site corresponding to nucleotides 303-304; and
 - (vii) the AU site corresponding to nucleotides 316-317.
- 29. A DNAzyme as claimed in claim 28 in which the cleavage site is the AU site corresponding to nucleotides 271-272.
- 30. A DNAzyme as claimed in claim 28 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
- 31. A DNAzyme as claimed in claim 29 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
- 32. A DNAzyme as claimed in claim 20 wherein each binding domain is nine or more nucleotides in length.
- 33. A DNAzyme as claimed in claim 32 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
- 34. A DNAzyme as claimed in claim 32 in which the cleavage site is selected from the group consisting of
 - (i) the GU site corresponding to nucleotides 198-199;
 - (ii) the GU site corresponding to nucleotides 200-201;
 - (iii) the GU site corresponding to nucleotides 264-265;
 - (iv) the AU site corresponding to nucleotides 271-272;
 - (v) the AU site corresponding to nucleotides 301-302;
 - (vi) the GU site corresponding to nucleotides 303-304; and
 - (vii) the AU site corresponding to nucleotides 316-317.
- 35. A DNAzyme as claimed in claim 34 in which the cleavage site is the AU site corresponding to nucleotides 271-272.
- 36. A DNAzyme as claimed in claim 34 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
- 37. A DNAzyme as claimed in claim 35 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
- 38. A DNAzyme as claimed in claim 32 in which the catalytic domain has the nucleotide sequence GGCTAGCTACAACGA [SEQ ID NO: 2].

- 39. A DNAzyme as claimed in claim 38 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
- 40. A DNAzyme as claimed in claim 38 in which the cleavage site is selected from the group consisting of
 - (i) the GU site corresponding to nucleotides 198-199;
 - (ii) the GU site corresponding to nucleotides 200-201;
 - (iii) the GU site corresponding to nucleotides 264-265;
 - (iv) the AU site corresponding to nucleotides 271-272;
 - (v) the AU site corresponding to nucleotides 301-302;
 - (vi) the GU site corresponding to nucleotides 303-304; and
 - (vii) the AU site corresponding to nucleotides 316-317.
- 41. A DNAzyme as claimed in claim 40 in which the cleavage site is the AU site corresponding to nucleotides 271-272.
- 42. A DNAzyme as claimed in claim 40 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
- 43. A DNAzyme as claimed in claim 41 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
- 44. A DNAzyme as claimed in claim 20 which has a sequence selected from the group consisting of:
 - (i) 5'-caggggacaGGCTAGCTACAACGAcgttgcggg (SEQ ID NO: 3);
 - (ii) 5'-tgcaggggaGGCTAGCTACAACGAaccgttgcg (SEQ ID NO: 4);
 - (iii) 5'-catcctggaGGCTAGCTACAACGAgagcagget (SEQ ID NO: 5);
 - (iv) 5'-ccgcggccaGGCTAGCTACAACGAcctggacga (SEQ ID NO: 6);
 - (v) 5'-ccgctgccaGGCTAGCTACAACGAcccggacgt (SEQ ID NO: 7);
 - (vi) 5'-gcggggacaGGCTAGCTACAACGAcagctgcat (SEQ ID NO: 8);
 - (vii) 5'-cagcggggaGGCTAGCTACAACGAatcagctgc (SEQ ID NO: 9); and
 - (viii) 5'-ggtcagagaGGCTAGCTACAACGActgcagcgg (SEQ ID NO: 10).
- 45. A DNAzyme as claimed in claim 44 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
- 46. A DNAzyme as claimed in claim 44 which has the sequence: 5'-ccgcggccaGGCTAGCTACAACGAcctggacga (SEQ ID NO: 6).
- 47. A DNAzyme as claimed in claim 46 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
- 48. A pharmaceutical composition comprising a DNAzyme according to any one of claims 20 to 47 and a pharmaceutically acceptable carrier.

- 49. A method of inhibiting EGR-1 activity in cells which comprises exposing the cell to a DNAzyme according to any one of claims 20 to 47.
- 50. A method as claimed in claim 49 wherein the cells are vascular cells.
- 51. A method as claimed in any one of claims 49 wherein the cells are cells involved in neoplasia.
- 52. A method of inhibiting proliferation or migration of cells in a subject which comprises administering to the subject a prophylactically effective dose of the pharmaceutical composition according to claim 48.
- 53. A method as claimed in claim 52 wherein the cells are vascular cells.
- 54. A method as claimed in any one of claims 52 wherein the cells are cells involved in neoplasia.
- 55. A method of treating a condition associated with cell proliferation or migration in a subject which comprises administering to the subject a therapeutically effective dose of the pharmaceutical composition according to claim 48.
- 56. A method as claimed in claim 55 wherein the cells are vascular cells.
- 57. A method as claimed in any one of claims 55 wherein the cells are cells involved in neoplasia.
- 58. A method as claimed in claim 55 wherein the condition associated with cell proliferation or migration is selected from the group consisting of post-angioplasty restenosis, vein graft failure, hypertension, transplant coronary disease, and complications associated with atherosclerosis or peripheral vascular disease.
- 59. An angioplastic stent for inhibition of the onset of restenosis, which comprises an angioplastic stent operably coated with a prophylactically effective dose of a DNAzyme according to any one of claims 20 to 47.
- 60. A method for inhibiting the onset of restenosis in a subject undergoing angioplasty, which comprises topically administering a prophylactically effective dose of a pharmaceutical composition according to claim 48 to the subject at around the time of the angioplasty.
- 61. A method according to claim 60 in which the pharmaceutical composition is administered by catheter.
- 62. A method for inhibiting the onset of restenosis in a subject undergoing angioplasty, which comprises topically administering a stent according to claim 58 to the subject at around the time of the angioplasty.

REMARKS

The Applicant is canceling the original claim set without prejudice to the Applicant's right to pursue claims of the same or similar scope in a duly filed continuing application, and is filing a new set of claims intended solely to place the claims in proper multiple dependent format. The following table demonstrates the new claims as they correspond to the original claims set:

New Claim No.	Original Claim No. and (Dependency)
Claim 20	Claim 1
Claim 21	Claim 8 (1)
Claim 22	Claim 4 (1)
Claim 23	Claim 5 (4)
Claim 24	Claim 8 (4)
Claim 25	Claim 8 (5)
Claim 26	Claim 3 (1)
Claim 27	Claim 8 (3)
Claim 28	Claim 4 (3)
Claim 29	Claim 5 (4)
Claim 30	Claim 8 (4)
Claim 31	Claim 8 (5)
Claim 32	Claim 2 (1)
Claim 33	Claim 8 (2)
Claim 34	Claim 4 (2)
Claim 35	Claim 5 (4)
Claim 36	Claim 8 (4)
Claim 37	Claim 8 (5)
Claim 38	Claim 3 (2)
Claim 39	Claim 8 (3)
Claim 40	Claim 4 (3)
Claim 41	Claim 5 (4)
Claim 42	Claim 8 (4)
Claim 43	Claim 8 (5)
Claim 44	Claim 6 (1)
Claim 45	Claim 8 (6)
Claim 46	Claim 7 (6)
Claim 47	Claim 8 (7)
Claim 48	Claim 9 (1-8)

New Claim No.	Original Claim No. and (Dependency)
Claim 49	Claim 10 (1-8)
Claim 50	Claim 13 (10)
Claim 51	Claim 14 (10)
Claim 52	Claim 11 (9)
Claim 53	Claim 13 (11)
Claim 54	Claim 14 (11)
Claim 55	Claim 12 (9)
Claim 56	Claim 13 (12)
Claim 57	Claim 14 (12)
Claim 58	Claim 15 (12)
Claim 59	Claim 16 (1-8)
Claim 60	Claim 17 (9)
Claim 61	Claim 18 (17)
Claim 62	Claim 19 (15)

The new claim set does not introduce new matter.

The abstract of the disclosure is identical to the abstract in the published

PCT pamphlet.

Respectfully submitted,

MARSHALL, O'TOOLE, GERSTEIN, MURRAY & BORUN

6300 Sears Tower

233 South Wacker Drive

Chicago, Illinois 60606

Dated: July 11, 2001

Joseph A. Williams, Jr. (Reg. No. 38,659) Attorney for Applicants

- 7 -

8/PR1S

JC18 Rec'd PCT/PTO 1 1 JUL 2001

WO 00/42173

5

10

15

20

25

30

PCT/AU00/00011

1

CATALYTIC MOLECULES

FIELD OF THE INVENTION

The present invention relates to DNAzymes which are targeted against mRNA molecules encoding EGR-1 (also known as Egr-1 or NGFI-A). The present invention also relates to compositions including these DNAzymes and to methods of treatment involving administration of the DNAzymes.

BACKGROUND OF THE INVENTION

Egr-1 expression in Smooth Muscle Cells

Smooth muscle cells (SMCs) are well recognized as a significant cellular component of atherosclerotic and post-angioplasty restenotic lesions (Stary et al, 1995; Holmes et al, 1984). SMC migration and proliferation are key events in the pathogenesis of these vascular disorders (Jackson & Schwartz, 1992; Libby et al, 1995). The promoter regions of many genes that encode mitogenic and migratory factors expressed by SMCs in these lesions (Evanko et al, 1998; Murry et al, 1996; Ueda et al, 1996; Tanizawa et al, 1996; Rekhter & Gordon, 1994; Hughes et al, 1993; Brogi et al, 1993; Wilcox et al 1989; Wilcox et al, 1988) contain nucleotide (nt) recognition elements for the nuclear protein and transcription factor, Egr-1 (Khachigian & Collins, 1997; Khachigian et al, 1996). Egr-1 is not expressed in the unmanipulated artery wall, but is rapidly activated by mechanical injury (Khachigian et al, 1996; Silverman et al, 1997; Kim et al, 1995). It is also induced in vascular endothelial cells and/or SMCs exposed to fluid biomechanical forces (Khachigian et al, 1997; Sumpio et al, 1998) and multiple other pathophysiologically-relevant agonists (Delbridge & Khachigian, 1997).

DNAzymes

In human gene therapy, antisense nucleic acid technology has been one of the major tools of choice to inactivate genes whose expression causes disease and is thus undesirable. The anti-sense approach employs a nucleic acid molecule that is complementary to, and thereby hybridizes with, an mRNA molecule encoding an undesirable gene. Such hybridization leads to the inhibition of gene expression.

10

15

20

25

30

35

PCT/AU00/00011

2

Anti-sense technology suffers from certain drawbacks. Anti-sense hybridization results in the formation of a DNA/target mRNA heteroduplex. This heteroduplex serves as a substrate for RNAse H-mediated degradation of the target mRNA component. Here, the DNA anti-sense molecule serves in a passive manner, in that it merely facilitates the required cleavage by endogenous RNAse H enzyme. This dependence on RNAse H confers limitations on the design of anti-sense molecules regarding their chemistry and ability to form stable heteroduplexes with their target mRNA's. Anti-sense DNA molecules also suffer from problems associated with non-specific activity and, at higher concentrations, even toxicity.

As an alternative to anti-sense molecules, catalytic nucleic acid molecules have shown promise as therapeutic agents for suppressing gene expression, and are widely discussed in the literature (Haseloff (1988); Breaker (1994); Koizumi (1989); Otsuka; Kashani-Sabet (1992); Raillard (1996); and Carmi (1996)). Thus, unlike a conventional anti-sense molecule, a catalytic nucleic acid molecule functions by actually cleaving its target mRNA molecule instead of merely binding to it. Catalytic nucleic acid molecules can only cleave a target nucleic acid sequence if that target sequence meets certain minimum requirements. The target sequence must be complementary to the hybridizing regions of the catalytic nucleic acid, and the target must contain a specific sequence at the site of cleavage.

Catalytic RNA molecules ("ribozymes") are well documented (Haseloff (1988); Symonds (1992); and Sun (1997)), and have been shown to be capable of cleaving both RNA (Haseloff (1988)) and DNA (Raillard (1996)) molecules. Indeed, the development of in vitro selection and evolution techniques has made it possible to obtain novel ribozymes against a known substrate, using either random variants of a known ribozyme or random-sequence RNA as a starting point (Pan (1992); Tsang (1994); and Breaker (1994)).

Ribozymes, however, are highly susceptible to enzymatic hydrolysis within the cells where they are intended to perform their function. This in turn limits their pharmaceutical applications.

Recently, a new class of catalytic molecules called "DNAzymes" was created (Breaker and Joyce (1995); Santoro (1997)). DNAzymes are single-stranded, and cleave both RNA (Breaker (1994); Santoro (1997)) and DNA (Carmi (1996)). A general model for the DNAzyme has been proposed, and is known as the "10-23" model. DNAzymes following the "10-23" model, also

15

20

25

30

35

3

referred to simply as "10-23 DNAzymes", have a catalytic domain of 15 deoxyribonucleotides, flanked by two substrate-recognition domains of seven to nine deoxyribonucleotides each. In vitro analyses show that this type of DNAzyme can effectively cleave its substrate RNA at purine:pyrimidine junctions under physiological conditions (Santoro (1997)).

DNAzymes show promise as therapeutic agents. However, DNAzyme success against a disease caused by the presence of a known mRNA molecule is not predictable. This unpredictability is due, in part, to two factors. First, certain mRNA secondary structures can impede a DNAzyme's ability to bind to and cleave its target mRNA. Second, the uptake of a DNAzyme by cells expressing the target mRNA may not be efficient enough to permit therapeutically meaningful results. For these reasons, merely knowing of a disease and its causative target mRNA sequence does not alone allow one to reasonably predict the therapeutic success of a DNAzyme against that target mRNA, absent an inventive step.

SUMMARY OF THE INVENTION

Accordingly, in a first aspect the present invention provides a DNAzyme which specifically cleaves EGR-1 mRNA, the DNAzyme including

- (i) a catalytic domain which cleaves mRNA at a purine:pyrimidine cleavage site;
- (ii) a first binding domain contiguous with the 5' end of the catalytic domain; and
- (iii) a second binding domain contiguous with the 3' end of the catalytic domain,

wherein the binding domains are sufficiently complementary to two regions immediately flanking a purine:pyrimidine cleavage site within the region of EGR-1 mRNA corresponding to nucleotides 168 to 332 as shown in SEQ ID NO:1, such that the DNAzyme cleaves the EGR-1 mRNA.

In a second aspect the present invention provides a pharmaceutical composition including a DNAzyme according to the first aspect and a pharmaceutically acceptable carrier.

In a third aspect the present invention provides a method of inhibiting EGR-1 activity in cells which includes exposing the cells to a DNAzyme according to the first aspect of the present invention.

5

10

15

PCT/AU00/00011

4

In a fourth aspect the present invention provides a method of inhibiting proliferation or migration of cells in a subject which includes administering to the subject a prophylactically effective dose of a DNAzyme according to the first aspect of the present invention.

In a fifth aspect the present invention provides a method of treating a condition associated with cell proliferation or migration in a subject which includes administering to the subject a prophylactically effective dose of a DNAzyme according to the first aspect of the present invention.

In a sixth aspect the present invention provides an angioplastic stent for inhibiting the onset of restenosis, which comprises an angioplastic stent operably coated with a prophylactically effective dose of a DNAzyme according to the first aspect.

In a seventh aspect, the present invention provides a method for inhibiting the onset of restenosis in a subject undergoing angioplasty, which comprises topically administering a stent according to the fifth aspect to the subject at around the time of the angioplasty.

PCT/AU00/00011

5

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 Sequence of NGFI-A DNAzyme (ED5), its scrambled control (ED5SCR) and 23 nt synthetic rat substrate. The translational start site is underlined.

Figure 2 NGFI-A DNAzyme inhibits the induction of NGFI-A mRNA and protein by serum. Northern blot analysis was performed with 25 μ g of total RNA. The blot was stripped and reprobed for β -Actin. Autoradiograms were analyzed by scanning densitometry and the ordinate axis is expressed as NGFI-A band intensity as a fraction of β -Actin band intensity. The mean and standard errors of the mean are indicated in the figure. Data is representative of 2 independent experiments. * indicates P<0.05 (Student's paired t-test) as compared to control (FBS alone).

15

20

25

10

Figure 3 SMC proliferation is inhibited by NGFI-A DNAzyme. a, Assessment of total cell numbers by Coulter counter. Growth-arrested SMCs that had been exposed to serum and/or DNAzyme for 3 days were trypsinized followed by quantitation of the suspension. The sequence of AS2 is 5'-CTT GGC CGC TGC CAT-3' (SEQ ID NO: 20). b, Proportion of cells incorporating Trypan Blue after exposure to serum and/or DNAzyme. Cells were stained incubated in 0.2% (w:v) Trypan Blue at 22 °C for 5 min prior to quantitation by hemocytometer in a blind manner. c, Effect of ED5 on pup SMC proliferation. Growth-arrested WKY12-22 cells exposed to serum and/or DNAzyme for 3 days were resuspended and numbers were quantitated by Coulter counter. Data is representative of 2 independent experiments performed in triplicate. The mean and standard errors of the mean are indicated in the figure. * indicates P<0.05 (Student's paired t-test) as compared to control (FBS alone).

30

35

Figure 4 NGFI-A DNAzyme inhibition of neointima formation in the rat carotid artery. Neointimal and medial areas of 5 consecutive sections per rat (5 rats per group) taken at 250 μ m intervals from the point of ligation were determined digitally and expressed as a ratio per group. The mean and standard errors of the mean are indicated by the ordinate axis. * denotes

PCT/AU00/00011

6

P<0.05 as compared to the Lig, Lig+Veh or Lig+Veh+ED5SCR groups using the Wilcoxen rank sum test for unpaired data. Lig denotes ligation, Veh denotes vehicle.

5 Figure 5 Selective inhibition of human smooth muscle cell proliferation by DzA.

Figure 6 Specific inhibition of porcine retinal smooth muscle cell proliferation by DzA.

10

15

20

25

30

35

PCT/AU00/00011

7

DETAILED DESCRIPTION OF THE INVENTION

Egr-1 (also known as NGFI-A and EGR-1) binds to the promoters of genes whose products influence cell movement and replication in the artery wall. Table 1 shows an alignment of the human EGR-1 cDNA sequence with the equivalent mouse (Egr-1) and rat (NGFI-A) sequences. The present inventors have now developed DNA-based enzymes that cut NGFI-A/Egr-1/EGR-1 RNA with high efficiency and specificity. The NGFI-A "DNAzyme" cleaved synthetic and in vitro transcribed NGFI-A RNA in a sequence-specific manner and inhibited production of NGFI-A in vascular smooth muscle cells without influencing levels of the related zinc finger protein, Sp1, or the immediate-early gene product, c-Fos. The DNAzyme blocked seruminducible DNA synthesis in smooth muscle cells and attenuated total cell proliferation. The DNAzyme also inhibited the reparative response to mechanical injury, both in culture and in the rat carotid artery wall. These results indicate a necessary and sufficient role for NGFI-A/Egr-1/EGR-1 in vascular smooth muscle cell growth and provide the first demonstration of a DNAzyme targeted against NGFI-A/Egr-1/EGR-1 transcripts.

Accordingly, in a first aspect the present invention provides a DNAzyme which specifically cleaves EGR-1 mRNA, the DNAzyme including

- (i) a catalytic domain which cleaves mRNA at a purine:pyrimidine cleavage site;
- (ii) a first binding domain contiguous with the 5' end of the catalytic domain; and
- (iii) a second binding domain contiguous with the 3' end of the catalytic domain,

wherein the binding domains are sufficiently complementary to two regions immediately flanking a purine:pyrimidine cleavage site within the region of EGR-1 mRNA corresponding to nucleotides 168 to 332 as shown in SEQ ID NO:1, such that the DNAzyme cleaves the EGR-1 mRNA.

As used herein, "DNAzyme" means a DNA molecule that specifically recognizes and cleaves a distinct target nucleic acid sequence, which may be either DNA or RNA

In a preferred embodiment of the first aspect of the present invention, the binding domains are complementary to the regions immediately flanking the cleavage site. It will be appreciated by those skilled in the art, however,

10

15

20

25

8

that strict complementarity may not be required for the DNAzyme to bind to and cleave the EGR-1 mRNA.

The catalytic domain of a DNAzyme of the present invention may be any suitable catalytic domain. Examples of suitable catalytic domains are described in *Santoro and Joyce*, 1997 and US 5807718, the entire contents of which are incorporated herein by reference. In a preferred embodiment, the catalytic domain has the nucleotide sequence GGCTAGCTACAACGA (SEQ ID NO: 2).

Within the parameters of the present invention, the binding domain lengths (also referred to herein as "arm lengths") can be of any permutation, and can be the same or different. In a preferred embodiment, the binding domain lengths are at least 6 nucleotides. Preferably, both binding domains have a combined total length of at least 14 nucleotides. Various permutations in the length of the two binding domains, such as 7+7, 8+8 and 9+9, are envisioned. It is well established that the greater the binding domain length, the more tightly it will bind to its complementary mRNA sequence. Accordingly, in a more preferred embodiment, each domain is nine or more nucleotides in length.

Within the context of the present invention, preferred cleavage sites within the region of EGR-1 mRNA corresponding to nucleotides 168 to 332 are as follows:

- (i) the GU site corresponding to nucleotides 198-199;
- (ii) the GU site corresponding to nucleotides 200-201;
- (iii) the GU site corresponding to nucleotides 264-265;
- (iv) the AU site corresponding to nucleotides 271-272;
- (v) the AU site corresponding to nucleotides 301-302;
- (vi) the GU site corresponding to nucleotides 303-304; and
- (vii) the AU site corresponding to nucleotides 316-317.

In a further preferred embodiment, the DNAzyme has a sequence selected from:

- (i) 5'-caggggacaGGCTAGCTACAACGAcgttgcggg (SEQ ID NO: 3) targets GU (nt 198, 199); arms hybridise to bp 189-207
- 35 (ii) 5'-tgcaggggaGGCTAGCTACAACGAaccgttgcg (SEQ ID NO: 4) targets GU (nt 200, 201); arms hybridise to bp 191-209

25

30

35

9

- (iii) 5'-catcctggaGGCTAGCTACAACGAgagcaggct (SEQ ID NO: 5) targets GU (nt 264, 265); arms hybridise to bp 255-273
- 5 (iv) 5'-ccgcggccaGGCTAGCTACAACGAcctggacga (SEQ ID NO: 6) targets AU (nt 271, 272); arms hybridise to bp 262-280
 - (v) 5'-ccgctgccaGGCTAGCTACAACGAcccggacgt (SEQ ID NO: 7) targets AU (nt 271, 272); arms hybridise to bp 262-280
 - (vi) 5'-gcggggacaGGCTAGCTACAACGAcagctgcat (SEQ ID NO: 8)targets AU (nt 301, 302); arms hybridise to bp 292-310
- (vii) 5'-cagcggggaGGCTAGCTACAACGAatcagctgc (SEQ ID NO: 9)
 targets GU (nt 303, 304); arms hybridise to bp 294-312
 - (viii) 5'-ggtcagagaGGCTAGCTACAACGActgcagcgg (SEQ ID NO: 10) targets AU (nt 316, 317); arms hybridise to bp 307-325.
- In a particularly preferred embodiment, the DNAzyme targets the AU site corresponding to nucleotides 271-272 (ie. the translation start codon).

In a further preferred embodiment, the DNAzyme has the sequence: 5'-ccgcggccaGGCTAGCTACAACGAcctggacga (SEQ ID NO: 6).

In applying DNAzyme-based treatments, it is preferable that the DNAzymes be as stable as possible against degradation in the intra-cellular milieu. One means of accomplishing this is by incorporating a 3'-3' inversion at one or more termini of the DNAzyme. More specifically, a 3'-3' inversion (also referred to herein simply as an "inversion") means the covalent phosphate bonding between the 3' carbons of the terminal nucleotide and its adjacent nucleotide. This type of bonding is opposed to the normal phosphate bonding between the 3' and 5' carbons of adjacent nucleotides, hence the term "inversion". Accordingly, in a preferred embodiment, the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3' end of the catalytic domain. In addition to inversions, the instant DNAzymes may contain modified nucleotides. Modified nucleotides

10

15

20

25

30

35

include, for example, N3'-P5' phosphoramidate linkages, and peptide-nucleic acid linkages. These are well known in the art.

In a particularly preferred embodiment, the DNAzyme includes an inverted T at the 3' position.

As will be appreciated by those skilled in the art, given that DNAzymes of the present invention cleave human EGR-1, similar DNAzymes can be produced to cleave the corresponding mRNA in other species, eg. rat (NGFI-A), mouse (Egr-1) etc. In a further aspect, the present invention provides such DNAzymes.

In a second aspect the present invention provides a pharmaceutical composition including a DNAzyme according to the first aspect and a pharmaceutically acceptable carrier.

In a third aspect the present invention provides a method of inhibiting EGR-1 activity in cells which includes exposing the cells to a DNAzyme according to the first aspect of the present invention.

In a fourth aspect the present invention provides a method of inhibiting proliferation or migration of cells in a subject which includes administering to the subject a prophylactically effective dose of a DNAzyme according to the first aspect of the present invention.

In a fifth aspect the present invention provides a method of treating a condition associated with cell proliferation or migration in a subject which includes administering to the subject a prophylactically effective dose of a DNAzyme according to the first aspect of the present invention.

In preferred embodiments of the third, fourth and fifth aspects of the present invention, the cells are vascular cells, particularly smooth muscle or endothelial cells. The cells may, however, be cells involved in neoplasia, such as tumour cells and the like.

Although the subject may be any animal or human, it is preferred that the subject is a human.

In a preferred embodiment, conditions associated with SMC proliferation(and migration) are selected from post-angioplasty restenosis, vein graft failure, transplant coronary disease and complications associated with atherosclerosis (cerebrovascular infarction (stroke), myocardial infarction (heart attack), hypertension or peripheral vascular disease (gangrene of the extremities).

Within the parameters of the fourth and fifth aspects of the present invention, any suitable mode of administration may be used to administer or deliver the DNAzyme.

In particular, delivery of the nucleic acid agents described may be achieved by one or more of the following methods:

5

10

15

20

25

30

35

- (a) Liposomes and liposome-protein conjugates and mixtures.
- (b) Using catheters to deliver intra-luminal formulations of the nucleic acid as a solution or in a complex with a liposome.
- (c) Catheter delivery to adventitial tissue as a solution or in a complex with a liposome.
- (d) Within a polymer formulation such as polyethylenimine (PEI) or pluronic gels or within ethylene vinyl acetate copolymer (EVAc). The polymer is preferably delivered intra-luminally.
- (e) The nucleic acid may be bound to a delivery agent such as a targetting moiety, or any suitable carrier such as a peptide or fatty acid molecule.
- (f) Within a viral-liposome complex, such as Sendai virus.
- (g) The nucleic acid may be delivered by a double angioplasty balloon device fixed to catheter.
- (h) The nucleic acid could be delivered on a specially prepared stent of the Schatz-Palmaz or derivative type. The stent could be coated with a polymer or agent impregnated with nucleic acid that allows controlled release of the molecules at the vessel wall.

In a preferred embodiment, the mode of administration is topical administration. Topical administration may be effected or performed using any of the various methods and delivery systems known to those skilled in the art. The topical administration can be performed, for example, via catheter and topical injection, and via coated stent as discussed below.

Pharmaceutical carriers for topical administration are well known in the art, as are methods for combining same with active agents to be delivered. The following delivery systems, which employ a number of routinely used carriers, are only representative of the many embodiments envisioned for administering the instant composition.

Topical delivery systems include, for example, gels and solutions, and can contain excipients such as solubilizers, permeation enhancers (e.g., fatty acids, fatty acid esters, fatty alcohols and amino acids), and hydrophilic

15

20

25

30

35

polymers (e.g., polycarbophil and polyvinylpyrolidone). In the preferred embodiment, the pharmaceutically acceptable carrier is a liposome or a biodegradable polymer. Examples of agents which can be used in this invention include the following: (1) CellFectin, 1:1.5 (M/M) liposome formulation of the cationic lipid N,N^I,N^{III},tetramethyl-N,N^I,N^{III},tetrapalmitylspermine and dioleoyl phosphatidyl-ethanolamine (DOPE) (GIBCO BRL); (2) Cytofection GSV, 2:1 (M/M) liposome formulation of a cationic lipid and DOPE (Glen Research); (3) DOTAP (N-[1-(2,3-dioleoyloxy)-N,N,N-trimethyl-ammoniummethylsulfate) (Boehringer Mannheim); (4) Lipofectamine, 3:1 (M/M) liposome formulation of the polycationic lipid DOSPA and the neutral lipid DOPE (GIBCO BRL); (5) FuGENE⁶ (Roche Molecular Biochemicals); (6) Superfect (Qiagen); and (7) Lipofectamine 2000 (Gibco-life Technologies).

Examples of suitable methods for topical administration of the DNAzymes of the present invention are described in Autieri et al. (1995), Simons et al. (1992), Morishita et al. (1993), Bennett and Schwartz (1995) and Frimerman et al. (1999).

Determining the prophylactically effective dose of the instant pharmaceutical composition can be done based on animal data using routine computational methods. In one embodiment, the prophylactically effective dose contains between about 0.1 mg and about 1 g of the instant DNAzyme. In another embodiment, the prophylactically effective dose contains between about 1 mg and about 100 mg of the instant DNAzyme. In a further embodiment, the prophylactically effective dose contains between about 10 mg and about 50 mg of the instant DNAzyme. In yet a further embodiment, the prophylactically effective dose contains about 25 mg of the instant DNAzyme.

In a sixth aspect the present invention provides an angioplastic stent for inhibiting the onset of restenosis, which comprises an angioplastic stent operably coated with a prophylactically effective dose of a DNAzyme according to the first aspect.

Angioplastic stents, also known by other terms such as "intravascular stents" or simple "stents", are well known in the art. They are routinely used to prevent vascular closure due to physical anomalies such as unwanted inward growth of vascular tissue due to surgical trauma. They often have a

WO 00/42173 PCT/AU00/00011

13

tubular, expanding lattice-type structure appropriate for their function, and can optionally be biodegradable.

In this invention, the stent can be operably coated with the instant pharmaceutical composition using any suitable means known in the art. Here, "operably coating" a stent means coating it in a way that permits the timely release of the pharmaceutical composition into the surrounding tissue to be treated once the coated stent is administered. Such coating methods, for example, can use the polymer polypyrrole.

In a seventh aspect, the present invention provides a method for inhibiting the onset of restenosis in a subject undergoing angioplasty, which comprises topically administering a stent according to the fifth aspect to the subject at around the time of the angioplasty.

10

15

As used herein, administration "at around the time of angioplasty" can be performed during the procedure, or immediately before or after the procedure. The administering can be performed according to known methods such as catheter delivery.

In order that the nature of the present invention may be more clearly understood, preferred forms thereof will now be described with reference to the following non-limiting Figures and Examples.

PCT/AU00/09011

14

Table 1

			Ta	able 1		
. 5	Symbol con	Gap	le: GenRunE Weight: 5.0 Weight: 0.3	00	ina.cmp Com	pCheck: 6876
	EGRlalıgn.		388 Type:		, 1998 12:0	7 Check: 5107
10	Name: mous Name: rath Name: huma		4388 Che	ck: 8587 W	Jeight: 1.0	SEQ ID NO:11) SEQ ID NO:12) (SEQ ID NO:1)
15	NB. THIS IS	RAT NGFI-A	numbering			50
15	mouseEgrl					
		CCGCGGAGCC	TCAGCTCTAC			GCGGGCGTCC
20		51				100
	mouseEGRl ratEGRl	CCGACTCCCG	CGCGCGTTCA	GGCTCCGGGT	TGGGAACCAA	GGAGGGGGAG
	humanEGR1					
25		101				150
23	mouseEGR1	101				130
	ratEGR1	GGTGGGTGCG	CCGACCCGGA	AACACCATAT	AAGGAGCAGG	AAGGATCCCC
	humanEGR1		• • • • • • • • • • • • • • • • • • • •		• • • • • • • • • • • • • • • • • • • •	
30		151				200
	mouseEGR1					
	ratEGR1 humanEGR1	CGCCGGAACA	GACCTTATTT			TGGCCCAATA
	HUMANDOKI					
35		201				250
	mouseEGR1 ratEGR1	TGGCCCTGCC	GCTTCCGGCT	CTEGENEGAG	GGGGGAACGG	GGGTTGGGGC
	humanEGR1			CIGGGAGGAG		
40						
40	mouseEGR1	251				300
	ratEGR1	GGGGGCAAGC	TGGGAACTCC		CCCGGGAGGC	
	humanEGR1					
45		301				350
	mouseEGR1					
	ratEGR1		TAGGCTTTCC	• • • • • • • • • • • • • • • • • • • •	GCGCTCAGGG	TGCCGGAGCC
	humanEGR1				• • • • • • • • •	•••••
50		351				400
	mouseEGR1 ratEGR1	CCTCCCACCC	TGGAAGCGCC	CACCCCTCTT	CCATCCCACC	 ፕሮሞሞር እርርሞር
	humanEGR1	GGTCGCAGGG				
55	mouseEGR1	401				450
	ratEGR1	ACTCCGGGTC	CTCCCGGTCG		ATTAGGGCTT	CCTGCTTCCC
	humanEGRl					
. 60		451				500

PCT/AU00/00011

	mouseEGR1 ratEGR1 humanEGR1		ATGTACGTCA		GGGCCCGTGC	TGTTTCAGAC
-						
5	mouseEGR1	501				550
	ratEGR1		GAGGCCGATT			
	humanEGRl	• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • •	CCGCAG
10		551				600
	mouseEGRl		GCCGCCGCCG	CGATTCGCCG	CCGCCGCCAG	
	ratEGR1		GCCGCCGCCG			
	humanEGR1	AACTTGGGGA	GCCGCCGCCG	CCATCCGCCG	CCGCAGCCAG	CTTCCGCCGC
15		601				650
	mouseEGR1		GCCCCTGCCC			
	ratEGR1 humanEGR1		GCCCCTGCCC			
		000110011000	3000013000	CAGCCTCCGC	AGCCGCGGCG	CGICCACGCC
20		651	m			700
	mouseEGR1 ratEGR1		TACCGCCAGC CACCGCCAGC			
	humanEGR1		CAGGGCGAGT			
25		707				
25	mouseEGRl	701	CCCGCATGTA	ACCCCCCCA A	CCCCCCCC	750
	ratEGR1		CCCGCATGTA			_
	humanEGR1		CCCGCATGTA			
30		751				800
-	mouseEGR1		GCCCCGGGCT	GCGCCCACC.	ACCCAACAT	
	ratEGRl		GCCCGGGCT			
	humanEGR1	TGCAGCTCCA	GCCCGGGCT	GCACCCCCC	GCCCGACAC	CAGCTCTCCA
35		801				850
	mouseEGR1	GCTCGCTGGT	CCGGGATGGC	AGCGGCCAAG	GCCGAGATGC	
	ratEGR1	GCTCGCACGT	CCGGGATGGC	AGCGGCCAAG	GCCGAGATGC	AATTGATGTC
	humanEGR1	GCCTGC <i>TCGT</i>	CCAGGATGGC	<i>CGCGG</i> CCAAG	GCCGAGATGC	AGCTGATGTC
40		ED5 (rat) a	rms hvbridi	se to bo 80	7-825 in ra	t secu
		hED5 (hum) a				
		851				900
	mouseEGR1		ATCTCTGACC	CGTTCGGCTC	CTTTCCTCAC	
45	ratEGR1		ATCTCTGACC			
	humanEGR1	CCCGCTGCAG	ATCTCTGACC	CGTTCGGATC	CTTTCCTCAC	TCGCCCACCA
		901				950
	mouseEGR1		CCCCAAACTG	GAGGAGATGA	TGCTGCTGAG	
50	ratEGR1	TGGACAACTA	CCCCAAACTG	GAGGAGATGA	TGCTGCTGAG	CAACGGGGCT
	humanEGR1	TGGACAACTA	CCCTAAGCTG	GAGGAGATGA	TGCTGCTGAG	CAACGGGGCT
		951				1000
	mouseEGR1	CCCCAGTTCC	TCGGTGCTGC	CGGAACCCCA	GAGGGCAGCG	GCGGTAAT
55	ratEGR1		TCGGTGCTGC			
	humanEGR1	CCCCAGTTCC	TCGGCGCCGC	CGGGGCCCCA	GAGGGCAGCG	GCAGCAACAG
		1001				1050
co	mouseEGRl		AGCAGCAGCA			
60	ratEGR1	CAGCAGCAGC	AGCAGCAGCA	GCAGCAGCGG	GGGCGGTGGT	GGGGGCGCA

PCT/AU00/00011

WO 00/42173

	humanEGR1	CAGCAGCAGC	AGCAGCGGGG	GCGGTGGAGG	CGGCGGGGC	GGCAGCAACA
		1051				1100
	mouseEGR1	GCAACAGCGG	CAGCAGCGCC	ТТСААТССТС	AAGGGGAGCC	
5	ratEGR1	GCAACAGCGG				
	humanEGRl	GCAGCAGCAG	CAGCAGCACC	TTCAACCCTC	AGGCGGACAC	GGGCGAGCAG
		1101				1150
10	mouseEGR1		ACCTGACCAC			
10	ratEGR1 humanEGR1		ACCTGACCAC ACCTGACCGC			
	Humanegri	CCCTACGAGC	ACCIGACCGC	AGAGICI	TTTCCTGACA	TCTCTCTGAA
		1151				1200
15	mouseEGR1		GCGATGGTGG			
15	ratEGR1		TGACTACCCT GTGCTGGTGG			
	humanEGR1	CAACGAGAAG	Greereeree	AGACCAGTTA	CCCCAGCCAA	ACCACTCGAC
		1201				1250
20	mouseEGR1		CACCTATACT			
20	ratEGR1		TAGGGACGGG			
	humanEGR1	TGCCCCCCAT	CACCTATACT	GGCCGCTTTT	CCCTGGAGCC	TGCACCCAAC
		1251				1300
0.5	mouseEGRl		CTTTGTGGCC			
25	ratEGR1		TGTGTTAGAG			
	humanEGR1	AGTGGCAACA	CCTTGTGGCC	CGAGCCCCTC	TTCAGCTTGG	TCAGTGGCCT
		1301				1350
30	mouseEGR1		ACCAATCCTC			
30	ratEGRI humanEGR1		CGCGGAGGGC			
	numanegar	AGIGAGCAIG	ACCAACCCAC	CGGCCTCCTC	GICCICAGCA	CCATCTCCAG
		1351				1400
3=	mouseEGR1		GTCTTCCTCT			
35	ratEGRI		GGCT			TTGTTTTGAT
	humanEGR1	CGGCCTCCTC	CGCCTCC	GCCTCCCAGA	GCCCACCCCI	GAGCIGCGCA
		1401				1450
40	mouseEGR1		ACGACAGCAG			
40	ratEGR1		GCCCCC			
	humanEGR1	GIGUCATUCA	ACGACAGCAG	TCCCATTTAC	TCAGCGGCAC	CCACCTTCCC
		1451				1500
4 55	mouseEGR1		ACTGACATTT			
45	ratEGR1		TCCCAAGGAA			
	humanEGRl	CACGCCGAAC	ACTGACATTT	TCCCTGAGCC	ACAAAGCCAG	GCCTTCCCGG
		1501				1550
	mouseEGR1		CACAGCCTTG			
50	ratEGRI		TAGGGGCGCG			
	humanEGRl	GCTCGGCAGG	GACAGCGCTC	CAGTACCCGC	CTCCTGCCTA	CCCTGCCGCC
		1551				1600
	mouseEGR1					TTCCACAACA
55	ratEGR1					GAGCTGCAGT
	humanEGR1	AAGGGTGGCT	TCCAGGTTCC	CATGATCCCC	GACTACCTGT	TTCCACAGCA
		1601				1650
	mouseEGRl					TTCCAGGGTC
60	ratEGR1	AGAGGGGGAT	TCTCTGTTTG	CGTCAGCTGT	CGAAATGGCT	CTGC

PCT/AU00/00011

WO 00/42173

	humanEGR1	GCAGGGGGAT	CTGGGCCTGG	GCACCCCAGA	CCAGAAGCCC	TTCCAGGGCC
- 5	mouseEGR1 ratEGR1 humanEGR1	CACTGGAGCA	TACCCAGCAG GGTCCAGGAA CACCCAGCAG	CATTGCAATC	TGCTGCTATC	AATTATTAAC
10	mouseEGR1 ratEGR1 humanEGR1	CACATCGAGA	CTCAGTCGGG GTCAGTGGTA CTCAGTCGGG	GCCGGGCGAC	CTCTTGCCTG	GCCGCTTCGG
15	mouseEGR1 ratEGR1 humanEGR1	CTCTCATCGT	CCAATCCCAG CCAGTGATTG CCAGTCCCAG	CTCTCCAGTA	ACCAGGCCTC	TCTGTTCTCT
20	mouseEGR1 ratEGR1 humanEGR1	TTCCTGCCAG	GCCCAGCAAG AGTCCTTTTC GCCCAGCAAG	TGACATCGCT	CTGAATAACG	AGAAGGCG
25	mouseEGR1 ratEGR1 humanEGR1	CTGGTGGAGA	CCTGCGATCG CAAGTTATCC CCTGTGATCG	CAGCCAAACT	ACCCGGTTGC	CTCCCATCAC
30	mouseEGR1 ·ratEGR1 humanEGR1	CTATACTGGC	ATCCACACAG CGCTTCTCCC ATCCACACAG	TGGAGCCTGC	ACCCAACAGT	GGCAACACTT
35	mcuseEGRl ratEGRl humanEGRl	TGTGGCCTGA	CAGTCGTAGT ACCCCTTTTC CAGCCGCAGC	AGCCTAGTCA	GTGGCCTTGT	GAGCATGACC
40	mouseEGRl ratEGRl humanEGRl	AACCCTCCAA	AGCCTTTTGC CCTCTTCATC AGCCCTTCGC	CTCAGCGCCT	TCTCCAGCTG	CTTCATCGTC
45	mouseEGRl ratEGRl humanEGRl	TTCCTCTGCC	CGCAAGAGGC TCCCAGAGCC CGCAAGAGGC	CACCCCTGAG	CTGTGCCGTG	CCGTCCAACG
50	mouseEGR1 ratEGR1 humanEGR1		CAAAAGTGTG CATTTACTCA CAAAAGTGTT	GCTGCACCCA	CCTTTCCTAC	TCCCAACACT
5 5	mouseEGR1 ratEGR1 humanEGR1		CCCCGGTTGC	GACATTTTTC	CTGAGCCCCA	AAGCCAGGCC
60	mouseEGR1 ratEGR1		CCTGCCACCA CTGCAGGCAC			2250 CCCACTTCCT CTGCCTACCC

PCT/AU00/00011

	humanEGR1	TTATCCATCC	CCGGCCACCA	CCTCATACCC	ATCCCCTGTG	CCCACCTCCT
5	mouseEGR1 ratEGR1 humanEGR1	TGCCACCAAG	TGGCTCCTCC GGTGGTTTCC CGGCTCCTCG	AGGTTCCCAT	GATCCCTGAC	TATCTGTTTC
10	mouseEGR1 ratEGRI humanEGR1	2301 CCGTCGCCGT CACAACAACA	CAGTGGCCAC GGGAGACCTG CGGTGGCCAC	CACCTTTGCC AGCCTGGGCA	TCCGTTCC`	2350 GAAGCCCTTC
15	mouseEGR1 ratEGR1 humanEGR1	2351 ACCTGC CAGGGTCTGG	TTTCCCCACC AGAACCGTAC TTTCCCGGCC	CAGGTCAGCA CCAGCAGCCT	GCTTCCCGTC TCGCTCACTC	2400 TGCGGGCGTC CACTATCCAC
20	mouseEGR1 ratEGR1 humanEGR1	TATCAAAGCC	TCAGCACCTC TTCGCCACTC TCAGCGCCTC	AGTCGGGCTC	CCAGGACTTA	AAGGCTCTTA
25	mouseEGR1 ratEGR1 humanEGR1	ATAACACCTA	ACAATTGAAA CCAGTCCCAA ACAATTGAAA	CTCATCAAAC	CCAGCCGCAT	GCGCAAGT
30	mouseEGRl ratEGRl humanEGRl	.ACCCCAÁCC	GGAGAGGCAG GGCCCAGCAA AAAGAAACAC	GACACCCCC	CATGAACGCC	CGTATGCTTG
35	mouseEGR1 ratEGR1 humanEGR1	CCCTGTTGAG	AAGAGGGGCC TCCTGCGATC AGGAGGGTT.	GCCGCTTTTC	TCGCTCGGAT	GAGCTTACAC
40	mouseEGR1 ratEGR1 humanEGR1	GCCACATCCG	CTACTCACGA CATCCATACA CCTCTCTACT	GGCCAGAA	GCCCTTCCAG	TGTCGAATCT
45	mouseEGRI ratEGR1 humanEGR1	GCATGCGTAA	TCCCTGCCTC TTTCAGTCGT TTCCCCTTCC	AGTGACCACC	TTACCACCCA	CATCCGCACC
50	mouseEGR1 ratEGR1 humanEGR1	CACACAGG	CATGTCCAAG CGAGAAGCCT CATGTCCAAG	TTTGCCTGTG	ACATTTGTGG	GAGAAAGTTT
55	mouseEGR1 ratEGR1 humanEGR1	GCCAGGAGTG	TATTGGAT ATGAACGCAA TTTTGGAT	GAGGCA:TACC	AAAATCCACT	TAAGACAGAA
60	mouseEGRl ratEGRl	2801 CCATC GGACAAGAAA	ACATGCCTGG GCAGACAAAA			

PCT/AU00/00011

	humanEGR1	CCATCA	TATGCCTGAC	CCCTTGCTCC	CTTCAATGCT	AGAAAATCGA
5	mouseEGR1 ratEGR1 humanEGR1	TCTCTTCCTA	AGAAAAAAA CCCATCCCCA	GTGGCTACCT	CCTACCCATC	CCCCGCCACC
10	mouseEGR1 ratEGR1 humanEGR1	ACCTCATTTC	CAGCATCTGT CATCCCCAGT CAGTGTCTGT	GCCCACCTCT	TACTCCTCTC	CGGGCTCCTC
15	mouseEGR1 ratEGR1 humanEGR1	TACCTACCCG	GATAATTTGC TCTCCTGCAC GATAATTTGC	ACAGTGGCTT	CCCATCGCCC	
20	mouseEGR1 ratEGR1 humanEGR1	CCACCTATGC	CTTTGGGG CTCCGTCC CTTGGGGGAA	CACCTGCTTT	CCCTGCCCAG	GTCAGCACCT
25	mouseEGRl ratEGRl humanEGRl	TCCAGTCTGC	ATTTTGTGAT AGGGGTCAGC ATTTTGTGAT	AACTCCTTCA	GCACCTCAAC	3100 GGGTCTTTCA
30	mouseEGR1 ratEGR1 humanEGR1		CATTTTTTT CAACCTTTTC ACCTTTTTTT	TCCTAGGACA	ATTGAAATTT	GCTAAAGGGA
35	mouseEGR1 ratEGR1 humanEGR1	ATGAAAGAGA	CAGAGTGTTG GCAAAGGGAG CAGAGTGTTG	GGGAGCGCGA	GAGACAATAA	AGGACAGGAG
40	mouseEGR1 ratEGR1 humanEGR1	.GGAAGAAAT	CACATGTGAC GGCCCGCAAG CACATGTGGC	AGGGGCTGCC	TCTTAGGTCA	GATGGAAGAT
45	mouseEGR1 ratEGR1 humanEGR1	CTCAGAGCCA	TTTTTGCCCG AGTCCTTCTA TTTTTCTTCG	GTCAGTAGAA	GGCCCGTTGG	CCACCAGCCC
50	mouseEGR1 ratEGR1 humanEGR1	TTTCACTTAG	TGTGACACGC CGTCCCTGCC TGTGATGCCC	CTC.CCCAGT	CCCGGTCCTT	TTGACTTCAG
55	mouseEGR1 ratEGR1 humanEGR1	CTGCCTGAAA	GGGACACGCT CAGCCACGTC GGGACATGCT	CAAGTTCTTC	$\mathtt{ACCT} \dots \mathtt{CTA}$	TCCAAAGGAC
60	mouseEGR1 ratEGR1		GGAGGCTTGA TGGTATTGGA			

PCT/AU00/00011

\G	G	GA	\GG	CT	TT	rG	GG	A GC	:AA	AAT	' A	AGG	AAC	SAGG		SCTG	AGCT	GΑ	
T	T	'GC	CTC	CCC	TT	C.	AG	CAC	CTA	GAA	C	ATC	:AA	STTG	(CTG.	35 TCTG AAAA TCTG	AA AA	
G	T	'GG	GGC	CCC	CTC	CA	GA.	ACC	CCT	GCC	C	TGI	'ATC	CTTT	' (STAC.	35 ATCT A ATAT		
Ġ	C.	AT	rge	JA1	CTI	ГТ	GT'	TT:	rcc	TTG	; G	GGI	TA	CTT	, (GATG	36 TTTT TGAA TTTT	GA	
ГC	Α	ACT	rci	rat	rT	GΤ	AC	TA!	rtt	GGZ	A G	TTF	'AA	rtci	٠ (CACT	36 TTGT TTGG TTGT	GG	
٩G	Α	AA	AG(CCF	AA(GC	AΑ	AC	CAA	TGO	3 T	'GAT	CC.	TCT#	١,	TTTT	37 TTGA GTGA TTGA	TG	
CA	G.	SA C	CAI	TT#	٩G	GT	ТT	GA.	AAC	TTT	rī	'TTI	TTT	TTTT	•	TGAA	37 GAAA GCAG AAAA	CA	
٩A	T	TT.	AA	CT	GG/	AG	CA	TG	TGI	'CA	3 <i>P</i>	GT	GTT	GTT	;	CGTT	38 LTAA	TT	
ÇG	C	CTC	CG	AC.	TG	TΑ	AC	TC	TCA	CA:	r	FTG/	ACA	AAA:	C .	ACGG	38 TTTC	STT	
ΤТ	T	TT	тт	GT'	TG	TT	TT	TG	AAA	\AA	A A	'AA'	TTT	TTT	Г	TTTC	39 50000	STC	
											T (GCC'	ттт	GTG	Г	GAC	39 ACAC	CTT	
AC			AC.	ΑТ	GT	GC	ΑP	TC	GT	GAG	G (CAC			ACCI	4 CTAC	000 gcc	
G.	 []	 AG	 ig.a	 GT	 GA	 TG	тт	 TC	AGG	 3GG	A (GGC	 TTT	 'AGA	G	CAC	4 GATG	050 AGG	

PCT/AU00/00011

	humanEGR1					
5	mouseEGR1 ratEGR1	4051 AAGAGGGCTG			CAATGTAAGA	
	humanEGR1			·····		
10	mouseEGR1 ratEGR1 humanEGR1	4101 AAAACAAAAA	TCTGAACTCT			
15	mouseEGR1 ratEGR1 humanEGR1	4151 GATTTATCCA	TGTTCGGGAG		GCGGTTACCT	
20	mouseEGR1 ratEGR1 humanEGR1	4201 CGGTGACTTT		GAACATGAAG		
25	mouseEGR1 ratEGR1 humanEGR1	4251 TTTTACTTCG		TGCTTAAACA		
30	mouseEGR1 ratEGR1 humanEGR1	4301 AAACACATTG		ACTGCCCATG		4350 TGTGTATCCT
35	mouseEGR1 ratEGR1 humanEGR1	4351 TCAGAAAAAT		ATAAAGAAAC		

5

10

15

20

25

30

35

PCT/AU00/00011

22

Example 1 Characterisation of DNAzymes ED5 and hED5

Materials and Methods

ODN synthesis. DNAzymes were synthesized commercially (Oligos Etc., Inc.) with an inverted T at the 3' position unless otherwise indicated. Substrates in cleavage reactions were synthesized with no such modification. Where indicated ODNs were 5'-end labeled with γ^{32} P-dATP and T4 polynucleotide kinase (New England Biolabs). Unincorporated label was separated from radiolabeled species by centrifugation on Chromaspin-10 columns (Clontech).

In vitro transcript and cleavage experiments. A ³²P-labelled 206 nt NGFI-A RNA transcript was prepared by in vitro transcription (T3 polymerase) of plasmid construct pJDM8 (as described in Milbrandt, 1987, the entire contents of which are incorporated herein by reference) previously cut with Bgl II. Reactions were performed in a total volume of 20 μl containing 10 mM MgCl₂, 5 mM Tris pH 7.5, 150 mM NaCl, 4.8 pmol of in vitro transcribed or synthetic RNA substrate and 60 pmol DNAzyme (1:12.5 substrate to DNAzyme ratio), unless otherwise indicated. Reactions were allowed to proceed at 37 °C for the times indicated and quenched by transferring an aliquot to tubes containing formamide loading buffer (Sambrook et al, 1989). Samples were run on 12% denaturing polyacrylamide gels and autoradiographed overnight at -80 °C.

Culture conditions and DNAzyme transfection. Primary rat aortic SMCs were obtained from Cell Applications, Inc., and grown in Waymouth's medium, pH 7.4, containing 10% fetal bovine serum (FBS), $50 \,\mu\text{g/ml}$ streptomycin and 50 IU/ml penicillin at 37 °C in a humidified atmosphere of 5% CO₂. SMCs were used in experiments between passages 3-7. Pup rat SMCs (WKY12-22 (as described in Lemire et al, 1994, the entire contents of which are incorporated herein by reference)) were grown under similar conditions. Subconfluent (60-70%) SMCs were incubated in serum-free medium (SFM) for 6 h prior to DNAzyme (or antisense ODN, where indicated) transfection (0.1 μ M) using Superfect in accordance with manufacturer's instructions (Qiagen). After 18 h, the cells were washed with phosphate-buffered saline (PBS). pH 7.4 prior to transfection a second time in 5% FBS.

15

20

25

30

35

Northern blot analysis. Total RNA was isolated using the TRIzol reagent (Life Technologies) and 25 μ g was resolved by electrophoresis prior to transfer to Hybond-N+ membranes (NEN-DuPont). Prehybridization, hybridization with α^{32} P-dCTP-labeled Egr-1 or β -Actin cDNA, and washing was performed essentially as previously described (Khachigian et al. 1995).

Western blot analysis. Growth-quiescent SMCs in 100 mm plates (Nunc-InterMed) were transfected with ED5 or ED5SCR as above, and incubated with 5% FBS for 1 h. The cells were washed in cold PBS, pH 7.4, and extracted in 150 mM NaCl, 50 mM Tris-HCl, pH 7.5, 1% sodium deoxycholate, 0.1% SDS, 1% Triton X-100, 5 mM EDTA, 1% trasylol, 10 µg/ml leupeptin, 1% aprotinin and 2 mM PMSF. Twenty four µg protein samples were loaded onto 10% denaturing SDS-polyacrylamide gels and electroblotted onto PVDF nylon membranes (NEN-DuPont). Membranes were air dried prior to blocking with non-fat skim milk powder in PBS containing 0.05% (w:v) Tween 20. Membranes were incubated with rabbit antibodies to Egr-1 or Sp1 (Santa Cruz Biotechnology, Inc.) (1:1000) then with HRP-linked mouse anti-rabbit Ig secondary antiserum (1:2000). Where mouse monoclonal c-Fos (Santa Cruz Biotechnology, Inc.) was used, detection was achieved with HRP-linked rabbit anti-mouse Ig. Proteins were visualized by chemiluminescent detection (NEN-DuPont).

Assays of cell proliferation. Growth-quiescent SMCs in 96-well titer plates (Nunc-InterMed) were transfected with ED5 or ED5SCR as above, then exposed to 5% FBS at 37 °C for 72 h. The cells were rinsed with PBS, pH 7.4, trypsinized and the suspension was quantitated using an automated Coulter counter.

Assessment of DNAzyme stability. DNAzymes were 5'-end labeled with γ^{32} P-dATP and separated from free label by centrifugation. Radiolabeled DNAzymes were incubated in 5% FBS or serum-free medium at 37 °C for the times indicated. Aliquots of the reaction were quenched by transfer to tubes containing formamide loading buffer (Sambrook et al, 1989). Samples were applied to 12% denaturing polyacrylamide gels and autoradiographed overnight at -80 °C.

SMC wounding assay. Confluent growth-quiescent SMCs in chamber slides (Nunc-InterMed) were exposed to ED5 or ED5SCR for 18 h prior to a single scrape with a sterile toothpick. Cells were treated with mitomycin C (Sigma) (20 μ M) for 2 h prior to injury (Pitsch et al, 1996; Horodyski &

15

20

25

30

35

24

Powell, 1996). Seventy-two h after injury, the cells were washed with PBS, pH 7.4, fixed with formaldehyde then stained with hematoxylin-eosin.

Rat arterial ligation model and analysis. Adult male Sprague Dawley rats weighing 300-350 g were anaesthetised using ketamine (60 mg/kg, i.p.) and xylazine (8 mg/kg, i.p.). The right common carotid artery was exposed up to the carotid bifurcation via a midline neck incision. Size 6/0 nonabsorbable suture was tied around the common carotid proximal to the bifurcation, ensuring cessation of blood flow distally. A 200 μ l solution at 4° C containing 500 μ g of DNAzyme (in DEPC-treated H₂O), 30 μ l of transfecting agent and Pluronic gel P127 (BASF) was applied around the vessel in each group of 5 rats, extending proximally from the ligature for 12-15 mm. These agents did not inhibit the solidification of the gel at 37 $^{\circ}$ C. After 3 days, vehicle with or without 500 $\mu \mathrm{g}$ of DNAzyme was administered a second time. Animals were sacrificed 18 days after ligation by lethal injection of phenobarbitone, and perfusion fixed using 10% (v:v) formaldehyde perfused at 120 mm Hg. Both carotids were then dissected free and placed in 10% formaldehyde, cut in 2 mm lengths and embedded in 3% (w:v) agarose prior to fixation in paraffin. Five μm sections were prepared at $250 \, \mu\mathrm{m}$ intervals along the vessel from the point of ligation and stained with hematoxylin and eosin. The neointimal and medial areas of 5 consecutive sections per rat were determined digitally using a customized software package (Magellan) (Halasz & Martin, 1984) and expressed as a mean ratio per group of 5 rats.

Results and Discussion

The 7x7 nt arms flanking the 15 nt DNAzyme catalytic domain in the original DNAzyme design 7 were extended by 2 nts per arm for improved specificity (L.-Q. Sun, data not shown) (Figure 1). The 3' terminus of the molecule was capped with an inverted 3'-3'-linked thymidine (T) to confer resistance to 3'->5' exonuclease digestion. The sequence in both arms of ED5 was scrambled (SCR) without altering the catalytic domain to produce DNAzyme ED5SCR (Figure 1).

A synthetic RNA substrate comprised of 23 nts, matching nts 805 to 827 of NGFI-A mRNA (Figure 1) was used to determine whether ED5 had the capacity to cleave target RNA. ED5 cleaved the ³²P-5'-end labeled 23-mer within 10 min. The 12-mer product corresponds to the length between the

10

PCT/AU00/00011

25

A(816)-U(817) junction and the 5' end of the substrate (Figure 1). In contrast, ED5SCR had no demonstrable effect on this synthetic substrate. Specific ED5 catalysis was further demonstrated by the inability of the human equivalent of this DNAzyme (hED5) to cleave the rat substrate over a wide range of stoichiometric ratios. Similar results were obtained using ED5SCR (data not shown): hED5 differs from the rat ED5 sequence by 3 of 18 nts in its hybridizing arms (Table 2). The catalytic effect of ED5 on a ³²P-labeled 206 nt fragment of native NGFI-A mRNA prepared by *in vitro* transcription was then determined. The cleavage reaction produced two radiolabeled species of 163 and 43 nt length consistent with DNAzyme cleavage at the A(816)-U(817) junction. In other experiments, ED5 also cleaved a ³²P-labeled NGFI-A transcript of 1960 nt length in a specific and time-dependent manner (data not shown).

Table 2. DNAzyme target sites in mRNA.

Similarity between the 18 nt arms of ED5 or hED5 and the mRNA of rat NGFI-A or human EGR-1 (among other transcription factors) is expressed as a percentage. The target sequence of ED5 in NGFI-A mRNA is 5'-807-A CGU CCG GGA UGG CAG CGG-825-3' (SEQ ID NO: 13) (rat NGFI-A sequence), and that of hED5 in EGR-1 is 5'-262-U CGU CCA GGA UGG CCG CGG-280-3' (SEQ ID NO: 14) (Human EGR-1 sequence). Nucleotides in bold indicate mismatches between rat and human sequences. Data obtained by a gap best fit search in ANGIS using sequences derived from Genbank and EMBL. Rat sequences for Sp1 and c-Fos have not been reported.

15	Gene	Accession number	Best homology over 18 nts (%)	
			ED5	hED5
	Rat NGFI-A	M18416	100	84.2
20	Human EGR-1	X52541	84.2	100
	Murine Sp1	AF022363	66.7	66.7
	Human c-Fos	K00650	66.7	66.7
	Murine c-Fos	X06769	61.1	66.7
	Human Sp1	AF044026	38.9	28.9
25				

25

30

35

To determine the effect of the DNAzymes on endogenous levels of NGFI-A mRNA, growth-quiescent SMCs were exposed to ED5 prior to stimulation with serum. Northern blot and densitometric analysis revealed that ED5 (0.1 μ M) inhibited serum-inducible steady-state NGFI-A mRNA levels by 55% (Figure 2a), whereas ED5SCR had no effect (Figure 2a). The capacity of ED5 to inhibit NGFI-A synthesis at the level of protein was assessed by Western blot analysis. Serum-induction of NGFI-A protein was suppressed by ED5. In contrast, neither ED5SCR nor EDC, a DNAzyme bearing an identical catalytic domain as ED5 and ED5SCR but flanked by nonsense arms had any influence on the induction of NGFI-A (data not

15

20

25

30

35

shown). ED5 failed to affect levels of the constitutively expressed, structurally -related zinc-finger protein, Sp1. It was also unable to block serum-induction of the immediate-early gene product, c-Fos whose induction, like NGFI-A, is dependent upon serum response elements in its promoter and phosphorylation mediated by extracellular-signal regulated kinase (Treisman, 1990, 1994 and 1995; Gashler & Sukhatme, 1995). These findings, taken together, demonstrate the capacity of ED5 to inhibit production of NGFI-A mRNA and protein in a gene-specific and sequence-specific manner, consistent with the lack of significant homology between its target site in NGFI-A mRNA and other mRNA (Table 2).

The effect of ED5 on SMC replication was then determined. Growth-quiescent SMCs were incubated with DNAzyme prior to exposure to serum and the assessment of cell numbers after 3 days. ED5 $(0.1\,\mu\text{M})$ inhibited SMC proliferation stimulated by serum by 70% (Figure 3a). In contrast, ED5SCR failed to influence SMC growth (Figure 3a). AS2, an antisense NGFI-A ODN able to inhibit SMC growth at $1\,\mu\text{M}$ failed to inhibit proliferation at the lower concentration (Figure 3a). Additional experiments revealed that ED5 also blocked serum-inducible ³H-thymidine incorporation into DNA (data not shown). ED5 inhibition was not a consequence of cell death since no change in morphology was observed, and the proportion of cells incorporating Trypan Blue in the presence of serum was not influenced by either DNAzyme (Figure 3b).

Cultured SMCs derived from the aortae of 2 week-old rats (WKY12-22) are morphologically and phenotypically similar to SMCs derived from the neointima of balloon-injured rat arteries (Seifert et al, 1984; Majesky et al, 1992). The epitheloid appearance of both WKY12-22 cells and neointimal cells contrasts with the elongated, bipolar nature of SMCs derived from normal quiescent media (Majesky et al, 1988). WKY12-22 cells grow more rapidly than medial SMCs and overexpress a large number of growth-regulatory molecules (Lemire et al, 1994), such as NGFI-A (Rafty & Khachigian, 1998), consistent with a "synthetic" phenotype (Majesky et al, 1992; Campbell & Campbell, 1985). ED5 attenuated serum-inducible WKY12-22 proliferation by approximately 75% (Figure 3c). ED5SCR had no inhibitory effect; surprisingly, it appeared to stimulate growth (Figure 3c). Trypan Blue exclusion revealed that DNAzyme inhibition was not a consequence of cytotoxicity (data not shown).

15

20

25

30

35

28

To ensure that differences in the biological effects of ED5 and ED5SCR were not the consequence of dissimilar intracellular localization, both DNAzymes were 5'-end labeled with fluorescein isothiocyanate (FITC) and incubated with SMCs. Fluorescence microscopy revealed that both FITC-ED5 and FITC-ED5SCR localized mainly within the nuclei. Punctate fluorescence in this cellular compartment was independent of DNAzyme sequence. Fluorescence was also observed in the cytoplasm, albeit with less intensity. Cultures not exposed to DNAzyme showed no evidence of autofluorescence.

Both molecules were 5'-end labeled with $\gamma^{32}P$ -dATP and incubated in culture medium to ascertain whether cellular responsiveness to ED5 and ED5SCR was a consequence of differences in DNAzyme stability. Both ^{32}P -ED5 and ^{32}P -ED5 are after 48 h. In contrast to ^{32}P -ED5 bearing the 3' inverted T, degradation of ^{32}P -ED5 bearing its 3' T in the correct orientation was observed as early as 1 h. Exposure to serum-free medium did not result in degradation of the molecule even after 48 h. These findings indicate that inverse orientation of the 3' base in the DNAzyme protects the molecule from nucleolytic cleavage by components in serum.

Physical trauma imparted to SMCs in culture results in outward migration from the wound edge and proliferation in the denuded zone. We determined whether ED5 could modulate this response to injury by exposing growth-quiescent SMCs to either DNazyme and Mitomycin C, an inhibitor of proliferation (Pitsch et al. 1996; Horodyski & Powell. 1996) prior to scraping. Cultures in which DNAzyme was absent repopulated the entire denuded zone within 3 days. ED5 inhibited this reparative response to injury and prevented additional growth in this area even after 6 days (data not shown). That ED5SCR had no effect in this system further demonstrates sequence-specific inhibition by ED5.

The effect of ED5 on neointima formation was investigated in a rat model. Complete ligation of the right common carotid artery proximal to the bifurcation results in migration of SMCs from the media to the intima where proliferation eventually leads to the formation of a neointima (Kumar & Lindner, 1997; Bhawan et al. 1977; Buck, 1961). Intimal thickening 18 days after ligation was inhibited 50% by ED5 (Figure 4). In contrast, neither its scrambled counterpart (Figure 4) nor the vehicle control (Figure 4) had any effect on neointima formation. These findings demonstrate the capacity of

ED5 to suppress SMC accumulation in the vascular lumen in a specific manner, and argue against inhibition as a mere consequence of a "mass effect" (Kitze et al, 1998; Tharlow et al, 1996).

Further experiments revealed the capacity of hED5 to cleave (human) EGR-1 RNA. hED5 cleaved its substrate in a dose-dependent manner over a wide range of stoichiometric ratios. hED5 also cleaved in a time-dependent manner, whereas hED5SCR, its scrambled counterpart, had no such catalytic property (data not shown).

The specific, growth-inhibitory properties of ED5 reported herein suggest that DNAzymes may be useful as therapeutic tools in the treatment of vascular disorders involving inappropriate SMC growth.

Example 2

Cleavage of human EGR-1 RNA by panel of candidate DNAzymes

15

20

30

17);

10

To evaluate which specific DNAzymes targeting human EGR-1 (other than hED5) efficiently cleave EGR-1 RNA, we prepared *in vitro* transcribed 35S-labeled EGR-1 RNA and incubated this substrate with candidate DNAzymes for various times. The EGR-1 plasmid template (hs164) was prepared by subcloning bps 168-332 of human EGR-1 into pGEM-T-easy. A 388 nt 35S-labeled substrate was prepared by *in vitro* transcription using SP6 polymerase. Time-dependent cleavage of the substrate was tested using the following DNZzymes:

DzA: 5'-CAGGGGACAGGCTAGCTACAACGACGTTGCGGG-X-3' (SEQ ID NO: 15);

DzB: 5'-TGCAGGGGAGGCTAGCTACAACGAACCGTTGCG-X-3' (SEQ ID NO: 16);

DzC: 5'-CATCCTGGAGGCTAGCTACAACGAGAGCAGGCT-X-3' (SEQ ID NO:

DzE: 5'-TCAGCTGCAGGCTAGCTACAACGACTCGGCCTT-X-3' (SEQ ID NO:

18); and
DzF: 5'-GCGGGGACAGGCTAGCTACAACGACAGCTGCAT-X-3' (SEQ ID NO: 19)

where X denotes a 3'-3-linked T.

10

15

25

35

PCT/AU00/00011

30

The DNAzyme that cleaved most effectively of this group was DzA, then DzB, then DzC. In contrast, DzE was inactive.

Example 3

5 Inhibition of induction of EGR-1 in human SMC by DzA

To determine whether DzA could block the induction of endogenous human EGR-1, we incubated growth-quiescent human aortic smooth muscle cells with 5% fetal bovine serum and observed the production of EGR-1 protein by Western blot analysis. This band representing the EGR-1 protein was blocked by 0.5 μ M DzA, delivered using FuGENE6 (Roche Molecular Biochemicals) and unaffected by DzE. The blot was then stripped and reprobed with antibodies to the transcription factor Spl. Results obtained showed that neither serum nor DzA affected induction of Sp1. A Coomassie Blue gel indicated that equal protein had been loaded.

The data demonstrate that DzA cleaves EGR-1 mRNA and blocks the induction of EGR-1 protein.

Example 4

20 Inhibition of human SMC proliferation by DzA

To ascertain whether proliferation of human SMCs could be inhibited by DzA, a population of SMCs was quantitated with and without exposure to DzA or DzE. SMC proliferation stimulated by 5% fetal bovine serum was significantly inhibited by 0.5 μ M DzA (Figure 5). In contrast, neither DzE nor ED5SCR had any effect (Figure 5). These data demonstrate that DzA inhibits human SMC proliferation.

Example 5

30 Inhibition of porcine SMC proliferation by DzA

The porcine and human EGR-1 sequences are remarkably well conserved (91%). Porcine retinal SMCs were used to determine whether DzA could block the growth of porcine SMCs. Our studies indicate that DzA (0.5 µM) could inhibit the proliferation of these cells (Figure 6). In contrast, DzE had no effect (Figure 6).

WO 60/42173 PCT/AU00/00011

31

Example 6 Delivery of DNAzyme into the porcine coronary artery wall

Porcine angioplasty and stenting are accepted models of human instent restenosis (Karas et al. 1992). The porcine coronary anatomy, dimensions and histological response to stenting are similar to the human (Muller et al. 1992). The Transport Catheter has previously been used to deliver antisense DNA targeting c-myc in humans (Serrys et al. 1998) and the pig (Gunn & Cumberland, 1996) via the intraluminal route. Using this catheter, FITC-labeled DNAzyme was applied to the inner wall of a porcine coronary artery, ex vivo, from a newly explanted pig heart. DNAzyme (1000 μg) was delivered via the catheter in 2ml MilliQ H20 containing 300μl FuGENE6 and 1mM MgC1₂. The FITC-labeled DNAzyme localised into the intimal cells of the vessel wall. These studies demonstrate that DNAzyme can be delivered to cells within the artery wall via an intraluminal catheter.

Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

20

25

30

It will be appreciated by persons skilled in the art that numerous variations and/or modifications may be made to the invention as shown in the specific embodiments without departing from the spirit or scope of the invention as broadly described. The present embodiments are, therefore, to be considered in all respects as illustrative and not restrictive. In addition, various documents are cited throughout this application. The disclosures of these documents are hereby incorporated by reference into this application to describe more fully the state of the art to which this invention pertains.

PCT/AU00/00011

32

References

Autieri, M.V. et al (1995) Antisense oligonucleotides to the P65 subunit of NF-κB inhibit human vascular smooth muscle cell adherence and proliferation and prevent neointiima formation in rat carotid arteries. Biochem; and Biophys. Res. Commun. 213:827-836.

Bennett, M.D. and Schwartz, S.M. (1995) Antisense Therapy for Angioplasty
10 Restenosis. Circulation 92:1981-1993.

Bhawan, J., Joris, I., DeGerolami, U. & Majno, G. (1977) Effect of occlusion on large vessels. Am. J. Pathol. 88, 355-380.

15 Breaker, R.R. and Joyce, G. (1994) Chemistry and Biology 1:223-229.

Breaker, R.R. and Joyce, G. (1995) Chemistry and Biology 2:655-660.

Brogi, E., et al. (1993) Distinct patterns of expression of fibroblast growth 20 factors and their receptors in human atheroma and nonatherosclerotic arteries. J. Clin. Invest. 92, 2408-2418.

Buck, R.C. (1961) Intimal thickening after ligature of arteries. Circ. Res. 9, 418-426.

Campbell, G.R. & Campbell, J.H. (1985) Smooth muscle phenotypic changes in arterial wall homeostasis: implications for the pathogenesis of atherosclerosis. Exp. Mol. Pathol. 42, 139-162.

30 Carmi, N., et al. (1996) Chemistry and Biology 3:1039-1046.

Delbridge, G.J. & Khachigian, L.M. (1997) FGF-1-induced PDGF A-chain gene expression in vascular endothelial cells involves transcriptional activation by Egr-1. Circ. Res. 81, 282-288.

25

10

20

35

Evanko, S.P., Raines, E.W., Ross, R., Gold, L.I. & Wight, T.N. (1998) Proteoglycan distribution in lesions of atherosclerosis depends on lesion severity, structural charcteristics, and the proximity of platelet-derived growth factor and transforming growth factor-beta. Am. J. Pathol. 152, 533-546.

Frimerman, A. et al. (1999) Chimeric DNA-RNA Hammerhead Ribozyme to Proliferating Cell Nuclear Antigen Reduces Stent-Induced Stenosis in a Porcine Coronary Model. Circulation 99: 697-703.

Gashler, A. & Sukhatme, V. (1995) Early growth response protein 1 (Egr-1): prototype of a zinc-finger family of transcription factors. Prog. Nucl. Acid Res. 50, 191-224.

Gunn J. and Cumberland D.C. (1996) Dual balloon catheter: seminars in Interventional Cardiology 1:31-33.

Halasz, P. & Martin, P. (1984) A microcomputer-based system for semi-automatic analysis of histological sections. Proc. Royal Microscop. Soc. 19, 312.

Haseloff, J. & Gerlach, W.A. (1988) Simple RNA enzymes with new and highly specific endoribonuclease activities. Nature 334, 585-591.

Holmes, D.R., et al. (1984) Restenosis after percutaneous transluminal coronary angioplasty (PTCA): a report from the PTCA Registry of the National Heart, Lung, and Blood Institute. Am. J. Cardiol. 53, 77C-81C.

Horodyski, J. & Powell, R.J. (1996) Effect of aprotinin on smooth muscle cell proliferation, migration, and extracellular matrix synthesis. J. Surg. Res. 66, 115-118.

Horodyski, J. & Powell, R.J. (1996) Effect of aprotinin on sincoth muscle cell proliferation, migration, and extracellular matrix synthesis. J. Surg. Res. 66, 115-118.

25

30

Hughes, S.E., Crossman, D. & Hall, P.A. (1993) Expression of basic and acidic fibroblast growth factors and their receptor in normal and atherosclerotic human arteries. Cardiovasc. Res. 27, 1214-1219.

Jackson, C.L. & Schwartz, S.M. (1992) Pharmacology of smooth muscle replication. Hypertension 20, 713-736.

Karas et al. (1992) Coronary intimal proliferation after balloon injury and stenting in the sine: an animal model of restenosis. *J. Am. Coll. Cardiol.* 20, 467-474.

Kashani-Sabet, M., et al. (1992) Antisense Research and Development 2:3-15.

15 Khachigian, L.M. & Collins, T. (1997) Inducible expression of Egr-1dependent genes: a paradigm of transcriptional activation in vascular endothelium. Circ. Res. 81, 457-461.

Khachigian, L.M., et al. (1997) Egr-1 is activated in endothelial cells exposed to fluid shear stress and interacts with a novel shear-stress response element in the PDGF A-chain promoter. Arterioscl. Thromb. Vasc. Biol. 17, 2280-2286.

Khachigian, L.M., Lindner, V., Williams, A.J. & Collins, T. (1996) Egr-1-induced endothelial gene expression: a common theme in vascular injury. Science 271, 1427-1431.

Khachigian, L.M., Williams, A.J. & Collins, T. (1995) Interplay of Sp1 and Egr-1 in the proximal PDGF-A promoter in cultured vascular endothelial cells. J. Biol. Chem. 270, 27679-27686.

Kim, S., et al. (1995) Angiotensin II type 1 receptor blockade inhibits the expression of immediate-early genes and fibronectin in rat injured artery. Circulation 92, 88-95.

Kitze, B., et al. (1998) Human CD4+ T lymphocytes recognize a highly conserved epitope of human T lymphotropic virus type 1 (HTLV-1) env gp21 restricted by HLA DRB1*0101. Clin. Exp. Immunol. 111, 278-285.

5 Koizumi, M., et al. (1989) Nucleic Acids Research 17:7059-7069.

Kumar, A. & Lindner, V. (1997) Remodeling with neointima formation in the mouse carotid artery after cessation of blood flow. Arterioscl. Thromb. Vasc. Biol. 17, 2238-2244.

10

Lemire, J.M., Covin, C.W., White, S., Giachelli, C.M. & Schwartz, S.M. (1994) Characterization of cloned aortic smooth muscle cells from young rats. Am. J. Pathol. 144, 1068-1081.

Libby, P., Schwartz, S.M., Brogi, E., Tanaka, H. & Clinton, S. (1995) A cascade model for restenosis. Circ. Res. 86 (Suppl. III), 47-52.

Majesky, M.W., Benditt, E.P. & Schwartz, S.M. (1988) Expression and developmental control of platelet-derived growth factor A-chain and B-chain/Sis genes in rat aortic smooth muscle cells. Proc. Natl. Acad. Sci. USA 85, 1524-1528.

Majesky, M.W., Giachelli, C.M., Reidy, M.A. & Schwartz, S.M. (1992) Rat carotid neointimal smooth muscle cells reexpress a developmentally regulated mRNA phenotype during repair of arterial injury. Circ. Res. 71, 759-768.

Milbrandt, J. (1987) A nerve growth factor-induced gene encodes a possible transcriptional regulatory factor. Science 238, 797-799

30

25

Morishita, R. et al (1993) Single intraluminal delivery of antisense cdc2 kinase and proliferating-cell nuclear antigen oligonucleotides result in chronic inhibition of neointimal hyperplasia. Proc. Natl. Acad. Sci. USA 90:8474-8478.

Muller DW et al. (1992) Experimental models of coronary artery restenosis. J. Am. Coll. Cardiol. 19, 418-432.

Murry, C.E., Bartosek, T., Giachelli, C.M., Alpers, C.E. & Schwartz, S.M. (1996) Platelet-derived growth factor-A mRNA expression in fetal, normal adult, and atherosclerotic human aortas. Circulation 93, 1095-1106

Otsuka, E. and Koizumi, M., Japanese Patent No. 4,235,919.

10 Pan, T. and Uhlenbeck, O.C. (1996) Biochemistry 31:3887-3895.

Pitsch, R.J., et al. (1996) Inhibition of smooth muscle cell proliferation and migration in vitro by antisense oligonucleotide to c-myb. J. Vasc. Surg. 23, 783-791.

Pitsch, R.J., et al. (1996) Inhibition of smooth muscle cell proliferation and migration in vitro by antisense oligonucleotide to c-myb. J. Vasc. Surg. 23, 783-791.

20 Raillard, S.A. and Joyce, G.F. (1996) Biochemistry 35:11693-11701.

Rafty, L.A. & Khachigian, L.M. (1998) Zinc finger transcription factors mediate high constitutive PDGF-B expression in smooth muscle cells derived from aortae of newborn rats. J. Biol. Chem. 273, 5758-5764.

Rekhter, M. & Gordon, D. (1994) Does platelet-derived growth factor-A chain stimulate proliferation of arterial mesenchymal cells in human atherosclerostic plaques? Circ. Res. 75, 410-417.

Sambrook, J., Fritsch, E.F. & Maniatis, T. (1989) Molecular cloning: a laboratory manual, Cold Spring Harbor Laboratory Press, Plainview, N.Y.

Santoro, S.W. & Joyce, G.F. (1997) A general purpose RNA-cleaving DNA enzyme. Proc. Natl. Acad. Sci. USA 94, 4262-4266.

15

25

Seifert, R.A., Schwartz, S.M. & Bowen-Pope, D.F. (1984) Developmentally regulated production of platelet-derived growth factor-like molecules. Nature 311, 669-671.

- 5 Serrys PW et al. (1998) Antisense oligonucleotide against c-myc administered with the Transport Catheter (36-48 holes, 250 µm diameter) for the prevention of in-stent restenosis: results of randomised ITALICS trial. Circulation: I1909.
- Silverman, E.S., Khachigian, L.M., Lindner, V., Williams, A.J. & Collins, T. (1997) Inducible PDGF A-chain transcription in vascular smooth muscle cells is mediated by Egr-1 displacement of Sp1 and Sp3. Am. J. Physiol. 42, H1415-H1426.
- Stary, H.C., et al. (1995) A definition of advanced types of atherosclerotic lesions and a histological classification of atherosclerosis: a report from the Committee on Vascular Lesions of the Council on Atherosclerosis, American Heart Association. Arterioscler. Thromb. Vasc. Biol. 15, 1512-1531.
- 20 Sumpio, B.E., et al. (1998) Regulation of PDGF-B by cyclic strain: lack of involvement of the shear-stress responsive element. Arterioscler. Thromb. Vasc. Biol. 18, 349-355.
 - Sun. L.Q., et al. (1997) Mol. Biotechnology 7:241-251.

Simons, M. et al (1992) Antisense c-myb oligonucleotides inhibit intimal arterial smooth muscle cell accumulation in vivo. Nature 359:67-70.

- Symonds, R.H. (1992) Ann. Rev. Biochem. 61:641-671.
- Tanizawa, S., Ueda, M., van der Loos, C.M., van der Wal, A.C. & Becker, A.E. (1996) Expression of platelet-derived growth factor B-chain and beta-receptor expression in human coronary arteries after percutaneous transluminal coronary angioplasty: an immunohisochemical study. Heart 75, 549-556

25

30

Tharlow, R.J., Hill, D.R. & Woodruff, G.N. (1996) Comparison of the autoradiographic binding distribution of [3H]-gabapentin with excitatory amino acid receptor and amino acid uptake site distributions in rat brain. Brit. J. Pharmacol. 118, 457-465.

5

Treisman, R. (1995) Journey to the surface of the cell: Fos regulation and the SRE. EMBO J. 14, 4905-4913.

Treisman, R. (1994) Ternary complex factor: growth factor regulated transcriptional activators. Curr. Opin. Genet. Develop. 4, 96-101.

Treisman, R. (1990) The SRE: a growth factor responsive transcriptional regulator. Sem. Cancer Biol. 1, 47-58.

15 Tsang, J. and Joyce, G.F. (1994) Biochemistry 33:5966-5973.

Ueda, M., et al. (1996) In situ detection of platelet-derived growth factor-A and -B chain mRNA in human coronary arteries after percutaneous transluminal coronary angioplasty. Am. J. Pathol. 149, 831-843.

20

Wilcox, J.N., Smith, K.M., Schwartz, S.M. & Gordon, D. (1989) Localization of tissue factor in the normal vessel wall and in the atherosclerotic plaque. Proc. Natl. Acad. Sci. USA 86, 2839-2843.

Wilcox, J.N., Smith, K.M., Williams, L.T., Schwartz, S.M. & Gordon, D. (1988)
Platelet-derived growth factor mRNA detection in human atherosclerotic
plaques by in situ hybridization. J. Clin. Invest. 82, 1134-1143.

15

25

30

39

Claims:

- 1. A DNAzyme which specifically cleaves EGR-1 mRNA, the DNAzyme comprising
 - (i) a catalytic domain which cleaves mRNA at a purine:pyrimidine cleavage site;
 - (ii) a first binding domain contiguous with the 5' end of the catalytic domain; and
- 10 (iii) a second binding domain contiguous with the 3' end of the catalytic domain,

wherein the binding domains are sufficiently complementary to the two regions immediately flanking a purine:pyrimidine cleavage site within the region of EGR-1 mRNA corresponding to nucleotides 168 to 332 as shown in SEQ ID NO:1, such that the DNAzyme cleaves the EGR-1 mRNA.

- 2. A DNAzyme as claimed in claim 1 wherein each binding domain is nine or more nucleotides in length.
- 20 3. A DNAzyme as claimed in claim 1 or claim 2 in which the catalytic domain has the nucleotide sequence GGCTAGCTACAACGA (SEQ ID NO: 2).
 - 4. A DNAzyme as claimed in any one of claims 1 to 3 in which the cleavage site is selected from the group consisting of
 - i) the GU site corresponding to nucleotides 198-199;
 - (ii) the GU site corresponding to nucleotides 200-201;
 - (iii) the GU site corresponding to nucleotides 264-265;
 - (iv) the AU site corresponding to nucleotides 271-272;
 - (v) the AU site corresponding to nucleotides 301-302;
 - (vi) the GU site corresponding to nucleotides 303-304; and
 - (vii) the AU site corresponding to nucleotides 316-317.
 - 5. A DNAzyme as claimed in claim 4 in which the cleavage site is the AU site corresponding to nucleotides 271-272.

30

35

40

- 6. A DNAzyme as claimed in claim 1 which has a sequence selected from the group consisting of:
- 5 (i) 5'-caggggacaGGCTAGCTACAACGAcgttgcggg (SEQ ID NO: 3);
 - (ii) 5'-tgcaggggaGGCTAGCTACAACGAaccgttgcg (SEQ ID NO: 4);
 - (iii) 5'-catcctggaGGCTAGCTACAACGAgagcagget (SEQ ID NO: 5);
 - (iv) 5'-ccgcggccaGGCTAGCTACAACGAcctggacga (SEQ ID NO: 6);
 - (v) 5'-ccgctgccaGGCTAGCTACAACGAcccggacgt (SEQ ID NO: 7);
- 10 (vi) 5'-gcggggacaGGCTAGCTACAACGAcagctgcat (SEQ ID NO: 8);
 - (vii) 5'-cagcggggaGGCTAGCTACAACGAatcagctgc (SEQ ID NO: 9); and
 - (viii) 5'-ggtcagagaGGCTAGCTACAACGActgcagcgg (SEQ ID NO: 10).
- 7. A DNAzyme as claimed in claim 6 which has the sequence: 5'-ccgcggccaGGCTAGCTACAACGAcctggacga (SEQ ID NO: 6).
 - 8. A DNAzyme as claimed in any one of claims 1 to 7, wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3' end of the catalytic domain.
 - 9. A pharmaceutical composition comprising a DNAzyme according to any one of claims 1 to 8 and a pharmaceutically acceptable carrier.
- 25 10. A method of inhibiting EGR-1 activity in cells which comprises exposing the cells to a DNAzyme according to any one of claims 1 to 8.
 - 11. A method of inhibiting proliferation or migration of cells in a subject which comprises administering to the subject a prophylactically effective dose of the pharmaceutical composition according to claim 9.
 - 12. A method of treating a condition associated with cell proliferation or migration in a subject which comprises administering to the subject a therapeutically effective dose of the pharmaceutical composition according to claim 9.

- 13. A method as claimed in any one of claims 10 to 12 wherein the cells are vascular cells.
- 5 14. A method as claimed in any one of claims 10 to 12 wherein the cells are cells involved in neoplasia.
 - 15. A method as claimed in claim 12 wherein the condition associated with cell proliferation or migration is selected from the group consisting of post-angioplasty restenosis, vein graft failure, hypertension, transplant coronary disease and complications associated with atherosclerosis or peripheral vascular disease.
- 16. An angioplastic stent for inhibiting the onset of restenosis, which
 15 comprises an angioplastic stent operably coated with a prophylactically effective dose of a DNAzyme according to any one of claims 1 to 8.
 - 17. A method for inhibiting the onset of restenosis in a subject undergoing angioplasty, which comprises topically administering a prophylactically effective dose of a pharmaceutical composition according to claim 9 to the subject at around the time of the angioplasty.
 - 18. A method according to claim 17 in which the pharmaceutical composition is administered by catheter.

25

20

10

19. A method for inhibiting the onset of restenosis in a subject undergoing angioplasty, which comprises topically administering a stent according to claim 15 to the subject at around the time of the angioplasty.

ABSTRACT

The present invention relates to DNAzymes which are targeted against mRNA molecules encoding EGR-1 (also known as Egr-1 and NGFI-A). The present invention also relates to compositions including these DNAzymes and to methods of treatment involving administration of the DNAzymes.

PCT/AU00/00011

1/8

Figure 1

ED5SCR

WO 00/42173 PCT/AU00/00011




Figure 2

PCT/AU00/00011




Figure 3A

PCT/AU00/00011




Figure 3B

PCT/AU00/00011




Figure 3C

PCT/AU00/00011




Figure 4

PCT/AU00/00011




Figure 5

PCT/AU00/00011




Figure 6



DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that my residence, post office address and citizenship are as stated below next to my name; believe that I am an original, first and joint inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled "CATALYTIC MOLECULES," the specification of which was filed as International Application No. PCT/AU00/00011 on January 11, 2000 and was transmitted to the U.S. Receiving Office on July 11, 2001 together with a Preliminary Amendment dated July 11, 2001, and which was assigned Application Serial No. 0.99/889.075. I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment(s) referred to above. I acknowledge the duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability as defined in 37 C.F.R. §1.56.

I hereby claim foreign priority benefits under 35 U.S.C. §119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:

			Priority Cl	Priority Claimed	
PP8103	Australia	11/January/1999	×		
(Application Serial Number)	(Country)	(Day/Month/Year Filed)	Yes	No	
I hereby claim the benefit	under 35 U.S.C. §119(e) of any Un	ited States provisional application(s) liste	d below:		

NONE
(Application Serial Number) (Day/Month/Year Filed)

I hereby claim the benefit under 35 U.S.C. §120 of any United States application(s) or PCT international application(s) designating the United States of America listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior application(s) in the manner provided by the first paragraph of 35 U.S.C. §112, I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in 37 C.F.R. §1.56 which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application:

NONE
(Application Serial Number) (Day/Month/Year Filed) (Status-Patented, Pending or Abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. §1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: I hereby appoint as my attorneys, with full powers of substitution and revocation, to prosecute this application and transact all business in the Patent and Trademark Office connected therewith:

24

John B. Lungmus(18,566)—Allen H. Gerstein (22,218)—Nate F. Scarpelli (22,320)—Michael F. Borun (25,447)—Trevor B. Joike (25,542)—Carl E. Moore, Jř. (26,487)

FIRM NAME

Richard H Anderson (26,526) Patrick D Ertel (26,877) Richard B Hoffman(26,910) James P Zeller (28,491) Kevin D Hogg (31,839) Jeffrey S Sharp (31,879)

PHONE NO.

Martin J. Hirsch (32,237)
James J. Napoli (32,361).
Richard M. La Barge (32,254).
Douglass C. Hochsteder (33,710).
Robert M. Gerstein (34,824).
Anthony G. Sitko (36,278)

CITY & STATE

James A. Flight (37,622).

Roger A Heppermann (37,641)

David A Gass (38,153)

Gregory C Mayer (38,238).

Michael R Weiner (38,359)

William K. Merkel (40,725)

ZIP CODE

Send correspondence to: David A. Gass

	wars Tower Wacker Drive Chicago, Illinois 60606-6402
Full Name of First or Sole Inventor David G. Atkins	Citizenship AU
Residence Address - Street 45W 60th St., Apt 17A	Post Office Address - Street 45W 60th St., Apt 17A
City (Zip) New York 10023	City (Zip) New York 10023
State or Country New York	State or Country New York
Date I NOVEMBER 2001	Signature // /////////////////////////////////

STREET

	
Second Joint Inventor, if any	Citizenship
Andrew R. Baker	AU
Residence Address - Street	Post Office Address - Street
3 Adelong Place	3 Adelong Place
City (Zip)	City (Zip)
Wahroonga 2076 WLS	Wahroonga 2076
State or Country	State or Country
New South Wales	New South Wales
Date 15 NOVEMBER 2501	Signature // //
≥ 12 Ngrand 12 50 1	

Third Joint Inventor, if any	Citizenship
Levon Michael Khachigian	AU
Residence Address - Street	Post Office Address - Street
5 Ratcliffe Street	5 Ratcliffe Street
City (Zip)	City (Zip)
Ryde 2112	Ryde 2112
State or Country	State or Country
New South Wales WLS	New South/Wales
Date 9 November 2001	Signature / luon Whalherien

APPLICABLE RULES AND STATUTES

37 CFR 1.56. DUTY OF DISCLOSURE - INFORMATION MATERIAL TO PATENTABILITY (Applicable Portion)

- (a) A patent by its very nature is affected with a public interest. The public interest is best served, and the most effective patent examination occurs when, at the time an application is being examined, the Office is aware of and evaluates the teachings of all information material to patentability. Each individual associated with the filing and prosecution of a patent application has a duty of candor and good faith in dealing with the Office, which includes a duty to disclose to the Office all information known to that individual to be material to patentability as defined in this section. The duty to disclose information exists with respect to each pending claim until the claim is canceled or withdrawn from consideration, or the application becomes abandoned. Information material to the patentability of a claim that is canceled or withdrawn from consideration need not be submitted if the information is not material to the patentability of any claim remaining under consideration in the application. There is no duty to submit information which is not material to the patentability of any existing claim. The duty to disclose all information known to be material to patentability of any claim issued in a patent was cited by the Office or submitted to the Office in the manner prescribed by §§ 1.97(b)-(d) and 1.98. However, no patent will be granted on an application in connection with which fraud on the Office was practiced or attempted or the duty of disclosure was violated through bad faith or intentional misconduct. The Office encourages applicants to carefully examine:
 - (1) prior art cited in search reports of a foreign patent office in a counterpart application, and
 - the closest information over which individuals associated with the filing or prosecution of a patent application believe any pending claim patentability defines, to make sure that any material information contained therein is disclosed to the Office.

Information relating to the following factual situations enumerated in 35 USC 102 and 103 may be considered material under 37 CFR 1.56(a).

35 U.S.C. 102. CONDITIONS FOR PATENTABILITY: NOVELTY AND LOSS OF RIGHT TO PATENT

A person shall be entitled to a patent unless --

- (a) the invention was known or used by others in this country, or patented or described in a printed publication in this or a foreign country, before the invention thereof by the applicant for patent, or
- (b) the invention was patented or described in a printed publication in this or a foreign country or in public use or on sale in this country, more than one year prior to the date of the application for patent in the United States, or
 - (c) he has abandoned the invention, or
- (d) the invention was first patented or caused to be patented, or was the subject of an inventor's certificate, by the applicant or his legal representatives or assigns in a foreign country prior to the date of the application for patent in this country on an application for patent or inventor's certificate filed more than twelve months before the filing of the application in the United States, or
- (e) the invention was described in a patent granted on an application for patent by another filed in the United States before the invention thereof by the applicant for patent, or on an international application by another who has fulfilled the requirements of paragraph (1), (2), and (4) of section 371(c) of this title before the invention thereof by the applicant for patent, or
 - (f) he did not himself invent the subject matter sought to be patented, or
- (g) before the applicant's invention thereof the invention was made in this country by another who had not abandoned, suppressed, or concealed it. In determining priority of invention there shall be considered not only the respective dates of conception and reduction to practice of the invention, but also the reasonable diligence of one who was first to conceive and last to reduce to practice, from a time prior to conception by the other.

35 U.S.C. 103. CONDITIONS FOR PATENTABILITY; NON-OBVIOUS SUBJECT MATTER (Applicable Portion)

A patent may not be obtained though the invention is not identically disclosed or described as set forth in section 102 of this title, if the differences between the subject matter sought to be patented and the prior art are such that the subject matter as a whole would have been obvious at the time the invention was made to a person having ordinary skill in the art to which said subject matter pertains. Patentability shall not be negatived by the manner in which the invention was made.

Subject matter developed by another person, which qualifies as prior art only under subsection (f) or (g) of section 102 of this title, shall not preclude patentability under this section where the subject matter and the claimed invention were, at the time the invention was made, owned by the same person or subject to an obligation of assignment to the same person.

35 U.S.C. 112. SPECIFICATION (Applicable Portion)

The specification shall contain a written description of the invention, and of the manner and process of making and using it, in such full, clear, concise, and exact terms as to enable any person skilled in the art to which it pertains, or with which it is most nearly connected, to make and use the same, and shall set forth the best mode contemplated by the inventor of carrying out his invention.

55

60

PCT/AU00/00011

1/8

```
SEQUENCE LISTING
        <110> Johnson & Johnson Pty Ltd
                   Unisearch Limited
 5
        <120> Catalytic molecules
        <150> PP 8103
        <151> 1999-01-11
10
        <160> 20
        <170> PatentIn Ver. 2.1
15
         <210> 1
         <211> 3132
         <212> DNA
        <213> Homo sapiens
20
         <400> 1
         cogoagaact tggggagoog cogoogooat cogoogoogo agocagotto ogoogoogoa 60
        ggaccggcc ctgcccagc ctccgcagcc gcggcgcgtc cacgcccgcc cgcgcccagg 120
         gegagteggg gtegeegeet geaegettet eagtgtteee egegeeeege atgtaaceeg 180
         gecaggeece egeaaeggtg teccetgeag etccageece gggetgeace ecceegeece 240
25
         gacaccaget etecageety etegteeagg atygeogegg ccaaggeega gatgeagetg 300
         atgteccege tgeagatete tgaecegtte ggateettte etcaetegee caecatggae 360
         aactacccta agctggagga gatgatgctg ctgagcaacg gggctcccca gttcctcggc 420
         ggaggeggeg ggggeggeag caacagcage ageageagea geacetteaa eceteaggeg 540
         gacacgggcg agcagecta cgageacetg accgeagagt etttteetga catetetetg 600
         aacaacgaga aggtgetggt ggagaccagt taccccagec aaaccaeteg actgeeeece 660
         atcacctata ctggccgctt ttccctggag cctgcaccca acagtggcaa caccttgtgg 720
         cccqaqccc tettcagett qqtcaqtqqc ctaqtqaqca tgaccaaccc accggcetec 780
         togtoctcag caccatotoc agoggootoc tecgeotocg ectoccagag eccacecetg 840
         agetgegeag tgecatecaa egaeageagt eccatttaet cageggeace cacetteece 900
35
         acgccgaaca ctgacatttt ccctgagcca caaagccagg ccttcccggg ctcggcaggg 960
         acagogotec agtacoogoc tectgeetac ectgeogoca agggtggett coaggtteec 1020
         atgatececq actacetott tecacageaq cagggggate tgggcetggg caccecagae 1080
         cagaageeet tecagggeet ggagageege acceageage ettegetaac ecetetgtet 1140
         actattaagg cctttgccac tcagtcgggc tcccaggacc tgaaggccct caataccagc 1200
40
         taccagtece ageteateaa acceageege atgegeaagt ateceaaceg geecageaag 1260
         acgeeeccc acgaacgee ttacgettge coagtggagt cetgtgateg ecgettetee 1320
         egeteegaeg ageteaceeg ceacateege atecacaeag gecagaagee ettecagtge 1380
         cgcatctgca tgcgcaactt cagccgcagc gaccacctca ccaccacat ccgcacccac 1440
         acaggegaaa ageeettege etgegacate tgtggaagaa agtttgecag gagegatgaa 1500
 45
         cgcáágággc ataccaagát ccácttgegg cágáággáca agaaagcagá caaaagtgtt 1560 gtggcetett eggceacete etetetet teetaceegt eeeeggttge tacetettae 1620
          cogtocoogg ttactaceto ttatocatoo coggeoacca cotoatacec atoccetgtg 1680
         occacctect tetectetee eggeteeteg acctacceat eccetgtgea cagtggette 1740
 50
         contrology cogregation can care to the control           agcagettee ettecteage tgteaccaae teetteageg eetecacagg gettteggae 1860
          atgacagcaa cottttotoo caggacaatt gaaatttgot aaagggaaag gggaaagaaa 1920
          gggaaaaggg agaaaaagaa acacaagaga cttaaaggac aggaggagga gatggccata 1980
          ggagaggagg gtteetetta ggteagatgg aggtteteag ageeaagtee teeeteteta 2040
```

ctggagtgga aggtetattg gecaacaate etttetgece aetteeeett eeccaattae 2100

tattcccttt gacttcaget geetgaaaca geeatgteea agttetteae etetateeaa 2160 agaacttgat ttgcatggat tttggataaa teattteagt ateateteea teatatgeet 2220 gacccettge teeetteaat getagaaaat egagttggea aaatggggtt tgggeeeete 2280 agageeetge eetgeaeeet tgtacagtgt etgtgeeatg gatttegttt ttettggggt 2340

actettgatg tgaagataat ttgcatatte tattgtatta tttggagtta ggteeteact 2400

PCT/AU00/00011

```
tgggggaaaa aaaaaaaaa aagccaagca aaccaatggt gatcctctat tttgtgatga 2460
    tgctqtqaca ataagtttga accttttttt ttgaaacagc agtcccagta ttctcagagc 2520
    atgtgtcaga gtgttgttcc gttaaccttt ttgtaaatac tgcttgaccg tactctcaca 2580
    tgtggcaaaa tatggtttgg tttttctttt tttttttga aagtgttttt tcttcgtcct 2640
    tttggtttaa aaagtttcac gtcttggtgc cttttgtgtg atgccccttg ctgatggctt 2700
    gacatgtgca attgtgaggg acatgctcac ctctagcctt aaggggggca gggagtgatg 2760
    agaatgtaag aaaacaaaat ctaaaacaaa atctgaactc tcaaaagtct attttttaa 2880
    ctgaaaatgt aaatttataa atatattcag gagttggaat gttgtagtta cctactgagt 2940
    aggoggogat tittgtatgt tatgaacatg cagttcatta tittgtggtt ctattttact 3000
    ttgtacttgt gtttgcttaa acaaagtgac tgtttggctt ataaacacat tgaatgcgct 3060
    ttattgccca tgggatatgt ggtgtatatc cttccaaaaa attaaaacga aaataaagta 3120
                                                                    3132
    gctgcgattg gg
15
    <210> 2
    <211> 15
    <212> DNA
     <213> Artificial Sequence
20
     <220>
     <223> Description of Artificial Sequence: Catalytic
          domain of DNAzyme
25
     <400> 2
                                                                    15
     ggctagctac aacga
     <210> 3
30
     <211> 33
     <212> DNA
     <213> Artificial Sequence
35
     <223> Description of Artificial Sequence: DNAzyme
                                                                     33
     caggggacag gctagctaca acgacgttgc ggg
40
     <210> 4
     <211> 33
     <212> DNA
     <213> Artificial Sequence
45
     <223> Description of Artificial Sequence: DNAzyme
                                                                     33
     tgcagggag gctagctaca acgaaccgtt gcg
50
     <210> 5
     <211> 33
55
     <212> DNA
     <213> Artificial Sequence
     <223> Description of Artificial Sequence: DNAzyme
60
```

WO 00/42173 PCT/AU00/00011 3/8

<400> 5 33 catcctggag gctagctaca acgagagcag gct 5 <210> 6 <211> 33 <212> DNA <213> Artificial Sequence 10 <223> Description of Artificial Sequence: DNAzyme 33 ccgcggccag gctagctaca acgacctgga cga 15 <210> 7 <211> 33 <212> DNA 20 <213> Artificial Sequence <223> Description of Artificial Sequence: DNAzyme 25 <400> 7 ccgctgccag gctagctaca acgacccgga cgt 33 <210> 8 30 <211> 33 <212> DNA <213> Artificial Sequence 35 <223> Description of Artificial Sequence: DNAzyme <400> 8 33 gcggggacag gctagctaca acgacagctg cat 40 <210> 9 <211> 33 <212> DNA <213> Artificial Sequence 45 <223> Description of Artificial Sequence: DNAzyme 33 50 cagcggggag gctagctaca acgaatcagc tgc <210> 10 <211> 33 55 <212> DNA <213> Artificial Sequence <223> Description of Artificial Sequence: DNAzyme 60

PCT/AU00/00011

4/8

<400> 10 33 ggtcagagag gctagctaca acgactgcag cgg <210> 11 <211> 3068 <212> DNA <213> Mus musculus 10 <400> 11 ggggageege cgeegegatt egeegeegee geeagettee geegeegeaa gateggeeee 60 tgccccagce teegeggeag ccetgegtee accaegggee geggetaceg ccageetggg 120 ggcccaceta cacteccege agtgtgccce tgcacecege atgtaacecg gccaacece 180 ggcgagtgtg ccctcagtag cttcggcccc gggctgcgcc caccacccaa catcagttct 240 15 ccageteget ggteegggat ggeageggee aaggeegaga tgeaattgat gteteegetg 300 cagatetetg accepting etectificet cacteaccea coatggacaa etaccecaaa 360 ctggaggaga tgatgctgct gagcaacggg gctccccagt tcctcggtgc tgccggaacc 420 ccagagggca gcggcggtaa tagcagcagc agcaccagca gcgggggcgg tggtggggc 480 ggcagcaaca gcggcagcag cgccttcaat cctcaagggg agccgagcga acaaccctat 540 gageacetga ecacagagte ettteetgae ategetetga ataatgagaa ggegatggtg 600 20 gagacgagtt atcccagcca aacgactcgg ttgcctccca tcacctatac tggccgcttc 660 tccctggagc ccgcacccaa cagtggcaac actttgtggc ctgaacccct tttcagccta 720 gtcagtggcc tcgtgagcat gaccaatcct ccgacctctt catcctcggc gccttctcca 780 gctgcttcat cgtcttcctc tgcctcccag agcccgccc tgagctgtgc cgtgccgtcc 840 aacgacagca gtcccatcta ctcggctgcg cccacctttc ctactcccaa cactgacatt 900 25 tttcctgagc cccaaagcca ggcctttcct ggctcggcag gcacagcctt gcagtacccg 960 cotcotgoot accotgooac caaaggtggt ttocaggtto coatgatece tgactatetg 1020 tttccacaac aacagggaga cctgagcctg ggcacccag accagaagcc cttccagggt 1080 ctggagaacc gtacccagca gccttcgctc actccactat ccactattaa agccttcgcc 1140 actcagtcgg gctcccagga cttaaaggct cttaatacca cctaccaatc ccagctcatc 1200 30 aaacccagcc gcatgcgcaa gtaccccaac cggcccagca agacaccccc ccatgaacgc 1260 ccatatgett geeetgiega giectgegat egeegetitt etegetegga tgagettaee 1320 cgccatatec gcatecacac aggccagaag ccettecagt gtcgaatetg catgcgtaac 1380 ttcagtcgta gtgaccacct taccacccac atccgcaccc acacaggcga gaagcetttt 1440 gestgtgaca tttgtgggag gaagtttgee aggagtgatg aacgeaagag geataceaaa 1500 35 atccatttaa gacagaagga caagaaagca gacaaaagtg tggtggcctc cccggctgcc 1560 tetteactet ettettace atceceagtg getacetect acceatece tgecaceace 1620 teatteceat eccetgtgee eacttectae tectetectg getectecae etacecatet 1680 cctgcgcaca gtggcttccc gtcgccgtca gtggccacca cctttgcctc cgttccacct 1740 gettteecea eccaggteag eagetteecg tetgegggeg teageagete etteageace 1800 teaactggte ttteagacat gacagegace tttteteeca ggacaattga aatttgetaa 1860 40 agggaataaa agaaagcaaa gggagaggca ggaaagacat aaaagcacag gagggaagag 1920 atggccgcaa gaggggccac ctcttaggte agatggaaga teteagagee aagteettet 1980 acteacgagt agaaggaceg ttggccaaca gccctttcac ttaccatccc tgcctccccc 2040 gtcctgttcc ctttgacttc agctgcctga aacagccatg tccaagttct tcacctctat 2100 ccaaaggact tgatttgcat ggtattggat aaatcatttc agtatcctct ccatcacatg 2160 45 cctggccctt gctcccttca gcgctagacc atcaagttgg cataaagaaa aaaaaatggg 2220 tttgggccct cagaaccctg ccctgcatct ttgtacagca tctgtgccat ggattttgtt 2280 ttccttgggg tattcttgat gtgaagataa tttgcatact ctattgtatt atttggagtt 2340 aaateeteae tttgggggag gggggagcaa ageeaageaa accaatgatg ateetetatt 2400 50 ttgtgatgac tctgctgtga cattaggttt gaagcatttt ttttttcaag cagcagtcct 2460 aggtattaac tggagcatgt gtcagagtgt tgttccgtta attttgtaaa tactggctcg 2520 actgtaactc tcacatgtga caaagtatgg tttgtttggt tgggttttgt ttttgagaat 2580 tittttgccc gtccctttgg tttcaaaagt ttcacgtctt ggtgcctttt gtgtgacacg 2640 ccttccgatg gcttgacatg cgcagatgtg agggacacgc tcaccttagc cttaaggggg 2700 55 tttcggtctc cagaatgtaa gaagaaaaaa tttaaacaaa aatctgaact ctcaaaaagtc 2820 tatttttttta aactgaaaat gtaaatttat acatctattc aggagttgga gtgttgtggt 2880 tacctactga gtaggctgca gtttttgtat gttatgaaca tgaagttcat tattttgtgg 2940 ttttatttta ctttgtactt gtgtttgctt aaacaaagta acctgtttgg cttataaaca 3000

60

PCT/AU00/00011

5/8

cattgaatgc gctctattgc ccatgggata tgtggtgtgt atccttcaga aaaattaaaa 3060 ggaaaaat <210> 12 <211> 4321 <212> DNA <213> Rattus rattus 10 <400> 12 ccqcqqaqcc tcaqctctac gcqcctqqcq ccctccctac gcgggcgtcc ccqactcccg 60 cgcgcgttca ggctccgggt tgggaaccaa ggaggggag ggtgggtgcg ccgacccgga 120 aacaccatat aaggagcagg aaggatcccc cgccggaaca gaccttattt gggcagcgcc 180 ttatatggag tggcccaata tggccctgcc gcttccggct ctgggaggag gggcgaacgg 240 gggttggggc gggggcaagc tgggaactcc aggagcctag cccgggaggc cactgccgct 300 gttccaatac taggetttcc aggageetga gegeteaggg tgccggagee ggtegeaggg 360 tggaagegee caeegetett ggatgggagg tetteaegte acteegggte etceeggteg 420 gteetteeat attagggett cetgetteee atatatggee atgtaegtea eggeggagge 480 gggcccgtgc tgtttcagac ccttgaaata gaggccgatt cggggagtcg cgagagatcc 540 cagegegeag aacttgggga geegeegeeg egattegeeg eegeegeeag etteegeege 600 egeaagateg geecetgee eageeteege ggeagecetg egteeaeeae gggeegegge 660 caccgccage ctgggggccc acctacacte cccgcagtgt gcccctgcac cccgcatgta 720 acceggecaa cateeggega gtgtgeeete agtagetteg geeeeggget gegeeeacea 780 cccaacatca getetecage tegeacgtee gggatggeag eggeeaagge egagatgeaa 840 25 ttgatgtctc cgctgcagat ctctgacccg ttcggctcct ttcctcactc acccaccatg 900 gacaactacc ccaaactgga ggagatgatg ctgctgagca acggggctcc ccagttcctc 960 ggtgctgccg gaaccccaga gggcagcggc ggcaataaca gcagcagcag cagcagcagc 1020 agcagogggg goggtggtgg gggcggcagc aacagoggca gcagogettt caatootcaa 1080 ggggagccga gcgaacaacc ctacgagcac ctgaccacag gtaagcggtg gtctgcgccg 1140 aggctgaatc ccccttcgtg actaccctaa cgtccagtcc tttgcagcac ggacctgcat 1200 30 ctagatetta gggaegggat tgggatttee etetatteea caeageteea gggaettgtg 1260 ttagagggat gtetggggac cocccaacce tocatecttg egggtgegeg gagggeagae 1320 cgtttgtttt ggatggagaa ctcaagttgc gtgggtggct ggagtgggg agggtttgtt 1380 ttgatgagea gggttgeece etececegeg egegttgteg egageettgt ttgeagettg 1440 ttcccaagga agggctgaaa tctgtcacca gggatgtccc gccgcccagg gtaggggcgc 1500 35 gcattagctg tggccactag ggtgctggcg ggattccctc accceggacg cctgctgcgg 1560 agogototoa gagotgoagt agagggggat tototgtttg ogtoagetgt ogaaatggot 1620 ctgccactgg agcaggtcca ggaacattgc aatctgctgc tatcaattat taaccacatc 1680 gagagteagt ggtageeggg egacetettg cetggeeget teggetetea tegteeagtg 1740 40 attgctctcc agtaaccagg cctctctgtt ctctttcctg ccagagtcct tttctgacat 1800 cgctctgaat aacgagaagg cgctggtgga gacaagttat cccagccaaa ctacccggtt 1860 geoteceate acetatactg geogettete ectggageet geacecaaca gtggcaacae 1920 tttgtggcct gaaccccttt tcagcctagt cagtggcctt gtgagcatga ccaaccctcc 1980 aacctettea tecteagege ettetecage tgetteateg tetteetetg ceteceagag 2040 occacecety agetytycey tyceytecaa cyacaycayt eccatttact cagetycace 2100 45 cacctttect acteceaaça etgacatttt teetgageee caaageeagg cettteetgg 2160 ctotgoaggo acagoottgo agtacoogoo tootgootac cotgocacca agggtggttt 2220 ccaggttccc atgatccctg actatctgtt tccacaacaa cagggagacc tgagcctggg 2280 cacccagac cagaageest tecagggtet ggagaacegt acccageage ettegeteae 2340 tocactatoc actatoaaag cottogocac toagtogggo toccaggact taaaggotot 2400 50 taataacacc taccagtccc aactcatcaa acccagccgc atgcgcaagt accccaaccg 2460 geocageaag acaeeeeee atgaacgeee gtatgettge cetgttgagt cetgegateg 2520 cogettetet egeteggatg agettacaeg ceacateege atecatacag gecagaagee 2580 cttccagtgt cgaatctgca tgcgtaattt cagtcgtagt gaccacctta ccaccacat 2640 cogcacccac acaggcgaga agccttttgc ctgtgacatt tgtgggagaa agtttgccag 2700 55 gagtgatgaa cgcaagaggc ataccaaaat ccacttaaga cagaaggaca agaaagcaga 2760 caaaagtgte gtggcetect cagetgeete tteeetetet teetacecat ceecagtgge 2820 tacctcctac ccatcccccg ccaccacctc atttccatcc ccagtgccca cctettactc 2880 ctctccqqqc tectetacci acceqtetec tqcacacaqt qqcttcccat cqccctcggt 2940

ggccaccacc tatgcctccg toccacctgc tttccctgcc caggtcagca ccttccagtc 3000

WO 00/42173 PCT/AU00/00011

6/8

```
tgcaggggtc agcaactcct tcagcacctc aacgggtctt tcagacatga cagcaacctt 3060
     tteteetagg acaattgaaa tttgetaaag ggaatgaaag agagcaaagg gaggggageg 3120 egagagacaa taaaggacag gagggaagaa atggeeegea agaggggetg cetettaggt 3180
     cagatggaag atctcagage caagteette tagtcagtag aaggeeegtt ggeeaccage 3240
     cetticaett agegiecetg ecetececag teceggiect titigaettea getgeetgaa 3300 acagecaegt ceaagitett cacetetate caaaggaett gattigeatg giattiggata 3360
     aaccatttea geatcatete caccacatge etggeeettg etecetteag cactagaaca 3420
     tcaagttggc tgaaaaaaa aatgggtctg ggccctcaga accctgccct gtatctttgt 3480
     acagcatctg tgccatggat tttgttttcc ttggggtatt cttgatgtga agataatttg 3540
     catactetat tgtactattt ggagttaaat teteaetttg ggggaggggg agcaaageca 3600 agcaaaccaa tggtgateet etattttgtg atgateetge tgtgacatta ggtttgaaac 3660
10
     ttttttttt ttttgaagca gcagtcctag gtattaactg gagcatgtgt cagagtgttg 3720 ttccgttaat tttgtaaata ctgctcgact gtaactctca catgtgacaa atacggttt 3780 gtttggttgg gttttttgtt gtttttgaaa aaaaaatttt tttttgccc gtccctttgg 3840
      tttcaaaagt ttcacgtctt ggtgcctttg tgtgacacac cttgccgatg gctggacatg 3900
15
     agaagaaaaa tttaaaaacaa aaatctgaac tctcaaaaagt ctatttttt aactgaaaat 4080
      gtagatttat ccatgttcgg gagttggaat gctgcggtta cctactgagt aggcggtgac 4140
      ttttgtatgc tatgaacatg aagttcatta ttttgtggtt ttattttact tcgtacttgt 4200
      gtttgcttaa acaaagtgac ttgtttggct tataaacaca ttgaatgcgc tttactgccc 4260
      atgggatatg tggtgtgtat ccttcagaaa aattaaaagg aaaataaaga aactaactgg 4320
                                                                                     4321
25
      <210> 13
      <211> 19
      <212> RNA
      <213> Rattus rattus
30
      <400> 13
                                                                                      19
      acguccggga uggcagcgg
35
      <210> 14
      <211> 19
      <212> RNA
      <213> Homo sapiens
40
      <400> 14
                                                                                      19
      ucquccagga uggccgcgg
      <210> 15
      <211> 34
 45
      <212> DNA
      <213> Artificial Sequence
      <220>
 50
       <221> misc_feature
       <222> (33)..(34)
       <223> 3'-3-linked T
       <223> Description of Artificial Sequence: DNAzyme
 55
                                                                                       34
       caggggacag gctagctaca acgacgttgc gggt
```

60

WO 00/42173 PCT/AU00/00011

```
<210> 16
     <211> 34
      <212> DNA
     <213> Artificial Sequence
      <220>
      <221> misc_feature
      <222> (33)..(34)
<223> 3'-3-linked T
10
      <220>
      <223> Description of Artificial Sequence: DNAzyme
      <400> 16
      tgcagggag gctagctaca acgaaccgtt gcgt
                                                                                    34
15
      <210> 17
      <211> 34
<212> DNA
20
      <213> Artificial Sequence
      <220>
      <221> misc_feature
      <222> (33)..(34)
<223> 3'-3-linked T
      <223> Description of Artificial Sequence: DNAzyme
30
                                                                                     34
      catcctggag gctagctaca acgagagcag gctt
35
     <210> 18
      <211> 34
<212> DNA
      <213> Artificial Sequence
     <220>
40
      <221> misc_feature
      <222> (33)..(34)
<223> 3'-3-linked T
45
      <220>
      <223> Description of Artificial Sequence: DNAzyme
                                                                                     34
      tcagctgcag gctagctaca acgactcggc cttt
50
      <210> 19
      <211> 34
<212> DNA
      <213> Artificial Sequence
      <220>
     <221> misc feature
<222> (33)..(34)
<223> 3'-3-linked T
 60
```

United States Patent & Trademark Office Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

Page(s) of TXPRESC MOW were not present for scanning.

Document title)

Page(s) of were not present for scanning.

Document title)

□ Scanned copy is best available.